

CHARACTERIZATION OF *BOTRYOCOCCUS SP.* AND IDENTIFICATION OF
CARBOHYDRATES METABOLIC RELATED ENZYMES TOWARD LIPID
PRODUCTION

ZUBAINATU ABBA

A Thesis Submitted in fulfilment for the Award of
Doctor of Philosophy in Science

Faculty of Applied Sciences and Technology
Universiti Tun Hussein Onn Malaysia

OCTOBER, 2019

I dedicate this work with the most profound sense of gratitude to Almighty Allah for giving me Strength and Inspiration in making my dreams come true.



ACKNOWLEDGEMENT

All Praises and thanks are to Allah Subhanahu Wa ta'ala for His infinite blessings and mercies. May the peace and blessings of Allah Subhanahu Wa ta'aala be upon Prophet Muhammad (Sallallahu alayhi Wa sallam).

I am grateful to Allah Subhanahu Wata'ala (SWT) that gave me the rare opportunity and sustained my life and health to carry out this research. My profound gratitude and appreciation goes to my friendly gentle sister supervisor, Dr Hazel Monica Matias- Peralta for your patience, cooperation and constant support in contributing immensely in giving guidance and all the assistance required for the progress of this research regardless of time and place

My sincere appreciation goes to my co-supervisor Dr Siti Fatima Muhammad Fauzi for her kindness, guidance and support throughout my study and my appreciation to my co supervisor Prof, Dr Yousif Abdalla Abakr from UNMC. I also wish to express my profound gratitude to Aisha Gogoba Ishaq for her timely intervention in seeing to the success of this work, thank you very much. Besides that, I would also like to thanks all my friends who gave me encouragement and support while carrying this research. Also, I would like to give thanks to everyone who had given me guidance and support during my study.

I am especially indebted to my parents and my brother Prof Mustafa Abba, who are and will always be there for me at every point of my life. Thank you for your ideal upbringing, love, sacrifices, and support. May Allah forgive them grant the Aljannatul Firdausi.

I want to give a special thanks to the Federal College of Education, Nigerian TETFUND and UTHM for giving me this opportunity to acquire this degree.

ABSTRACT

The advancement of microalgal biofuels is faced with challenges among which is low strain performance in lipid production. Alternatively, the carbohydrate production in microalgal cells could be manipulated to increase the lipid content thereby maximizing the overall biofuel production. This work is aimed at determining metabolic pathway and enzymes involved in carbohydrate metabolism of *Botryococcus sp.* The *B. sp.* from Taman Negara Johor Endau Rompin was isolated and identified. Some relevant intracellular metabolites were extracted and quantified through HPLC and GC-MS analysis while extracellular metabolites excreted into the Bold basal medium were also analysed and identified through GC-MS. Enzymes involved in carbohydrate metabolism leading to lipid production in *B. sp.* under natural conditions were also identified through one-dimensional gel electrophoresis followed by proteomic mass spectrometry (LC-MS) and database searching. Finally, the carbohydrate to lipid metabolic pathway of *B. sp.* cultivated under natural conditions was determined. The total carbohydrate content was found to be 23 % per milligram biomass dry weight with monomeric sugars galactose, glucose, mannose, and arabinose. Total protein estimated for the microalgae *B. sp.* is 16.22 % and the lipid content was found to be 60.69 %. The extracellular metabolites constitute majorly cyclohydrocarbons, nitrogenated hydrocarbons, siloxanes, phenols and phenol derivatives. A glycolytic enzyme 'Enolase' which can generate phosphoenolpyruvate (PEP) and then convert it into pyruvate was identified in this study. Enolase which helps in high lipid metabolism was found in the cytoplasm and was used to construct the alternative pathway. Enolase was found to export fixed carbon (3PGA) to the cytoplasm, hence providing a shorter route to lipid production than the normal process via the plastid leading to the production of more lipid in *Botryococcus*.

ABSTRAK

Kemajuan biofuels biojisim berhadapan dengan cabaran di mana prestasi strain adalah rendah dalam pengeluaran lipid. Selain itu, pengeluaran karbohidrat dalam sel-sel mikroalga boleh dimanipulasi untuk meningkatkan kandungan lipid, seterusnya memaksimumkan pengeluaran biofuel secara menyeluruh. Kajian ini bertujuan untuk menentukan laluan metabolik dan enzim yang terlibat dalam metabolisme karbohidrat *Botryococcus sp. B. sp.* dari Taman Negara Johor Endau Rompin diasing dan dikenalpasti. Sesetengah metabolit intraselular yang berkaitan telah diekstrak dan dikira melalui analisis HPLC dan GC-MS manakala metabolit ekstraselular yang dikumuhkan ke dalam medium basal Bold juga dianalisis dan dikenalpasti melalui GC-MS. Enzim yang terlibat dalam metabolisme karbohidrat yang membawa kepada pengeluaran lipid dalam *B. sp.* dibawah keadaan semulajadi juga dikenalpasti melalui elektroforesis gel satu dimensi diikuti dengan spektrometri jisim proteomik (LC-MS) dan pencarian pangkalan data. Akhirnya, laluan metabolik karbohidrat ke lipid *B. sp.* terkultur di bawah keadaan semulajadi telah ditentukan. Jumlah kandungan karbohidrat didapati 23 % per miligram berat kering biomas dengan gula monomerik galaktosa, glukosa, mannososa, dan arabinosa. Jumlah protein yang dianggarkan untuk *B. sp.* mikroalga adalah 16.22 % dan kandungan lipid didapati 60.69 %. Metabolit ekstraselular merupakan sebahagian besar daripada siklohidrokarbon, hidrokarbon nitrogenasi, siloxan, fenol dan terbitan fenol. Enzim glikolitik 'Enolase' yang boleh menghasilkan fosfoenolpyruvat (PEP) dan kemudian mengubahnya menjadi piruvat telah dikenalpasti dalam kajian ini. Enolase yang membantu dalam metabolisme lipid tinggi didapati di sitoplasma dan digunakan untuk membina laluan alternatif. Enolase ditemui untuk mengeksport karbon tetap (3PGA) ke sitoplasma, dengan itu memberikan laluan yang lebih pendek ke pengeluaran lipid daripada proses biasa melalui plastid yang membawa kepada pengeluaran lebih banyak lipid dalam *Botryococcus*.

TABLES OF CONTENTS

TITLE PAGE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLES OF CONTENTS	vii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
LIST OF APPENDICIES	xviii
CHAPTER 1 INTRODUCTION	1
1.1 Background	1
1.2 Statement of the problem	3
1.3 Research aim and objectives	4
1.4 The importance of the study	5
1.5 The scope of the study	6
CHAPTER 2 LITERATURE REVIEW	7
2.1 Microalgae, <i>Botryococcus</i> sp.	7
2.2 Factors influencing the growth of microalgae	9
2.2.1 Photoperiod	9
2.2.2 Light	9

2.2.3	Temperature	10
2.2.4	pH	11
2.2.5	Mixing	12
2.2.6	Carbon	13
2.2.7	Nitrogen	13
2.2.8	Phosphorous	14
2.2.9	Micronutrients	14
2.3	Photosynthesis	15
2.3.1	Carboxylative phase:	16
2.3.2	Reductive phase	17
2.3.3	Regenerative phase	18
2.4	Thermogravimetric analysis (TGA)	19
2.5	Metabolomics	20
2.5.1	Intracellular Metabolites	21
2.5.2	Excretory products from microalgae (Extracellular metabolites)	23
2.6	Proteomics	25
2.6.1	Types of proteomics	25
2.6.2	Separation of proteins	26
2.6.3	Identification of proteins	27
2.7	Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS PAGE)	28
2.7.1	Basic Local Alignment Search Tool (BLAST)	29
2.7.2	BRENDA	30
2.8	Microalgal Enzymes	31
2.8.1	Cellulases	31

2.8.2	Enolase	32
2.8.3	Galactosidases	34
2.8.4	Carbohydrate composition of microalgae	35
2.8.5	Carbohydrate metabolism pathway of microalgae	36
2.8.6	Enzymes involved in carbohydrate accumulation	40
2.9	Manipulation of metabolic pathways of lipid accumulation	40
2.10	Lipid biosynthesis.	41
2.11	Mass production of microalgae	43
2.11.1	Manipulation of biomass production in a photobioreactors	44
2.11.2	Temperature	45
2.11.3	Light intensity	46
2.11.4	Nutrients starvation	47
2.11.5	Carbon dioxide	48
2.12	Biomass as a source of biofuel	49
2.13	Conclusion	50
CHAPTER 3	METHODOLOGY	54
3.1	Introduction	54
3.2	Materials	54
3.3	Equipment's	55
3.4	Microalgae	55
3.5	Flow of research	56
3.6	Microalgal medium preparation	57
3.6.1	Preparation of bold basal medium	

broth (BBM)	57
3.6.2 The subculture of <i>Botryococcus sp.</i> in the lab	58
3.6.3 Determination of microalgal cell density	58
3.6.4 Harvesting	59
3.6.5 Freeze-drying	59
3.7 Characterisation and quantification of the elemental composition of microalgal cells of <i>Botryococcus sp.</i>	59
3.8 Proximate analysis	60
3.9 Determination of Total Carbohydrates	60
3.9.1 Preparation of calibration standard	60
3.9.2 Hydrolysis of <i>Botryococcus sp.</i> biomass	61
3.9.3 HPLC analysis	61
3.10 Total protein	61
3.10.1 Protein extraction	61
3.10.2 Protein assay by Lowry method	62
3.10.3 Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS- PAGE) for Protein Separation	62
3.10.4 Sample preparation	63
3.10.5 Electrophoresis	64
3.10.6 Staining and destaining of gel	65
3.11 Sample preparation for protein identification from the Polyacrylamide gel for mass spectrometric analysis (LC-MS)	66
3.11.1 Reduction of disulfide bonds and alkylate free cysteine's	66

3.11.2	Enzymatic digestion	67
3.11.3	Extraction of peptides	67
3.12	Lipid extraction	67
3.12.1	Preparation for the extraction	68
3.12.2	Sample preparation	68
3.12.3	Soxhlet extraction procedure	68
3.13	Total fatty acids analysis	68
CHAPTER 4	RESULTS AND DISCUSSION	69
4.1	Production of <i>Botryococcus</i> biomass	69
4.2	Morphological and molecular identification of <i>Botryococcus</i> sp. cells	70
4.2.1	Quantification of the elemental composition of microalgal cells of <i>Botryococcus</i> sp. using Energy Dispersive X-ray spectroscopy	72
4.3	Proximate analysis	74
4.3.1	Thermogravimetric analysis	75
4.4	Intracellular metabolites of <i>Botryococcus</i> sp	77
4.4.1	Total sugars content	77
4.5	Total Protein estimation	79
4.5.1	Total Lipid estimation and fatty acid analysis	81
4.6	Extracellular metabolites	83
4.6.1	Carbohydrate metabolism pathway of <i>Botryococcus</i> sp.	87
4.6.2	Alternative pathway	92
4.6.3	Blast protein results	93

CHAPTER 5 CONCLUSION AND RECOMMENDATIONS	95
5.1 Conclusion	95
5.2 Recommendations	95
REFERENCES	97
APPENDICES	131
VITA	



LIST OF TABLES

2.1	Composition of microalgal cell wall and storage products. (Adapted from (Chen <i>et al.</i> , 2013)	36
2.2	Summarised literature	52
3.1	Composition of bold basal medium (BBM) (Nichols, 1973; Nichols & Bold, 1965)	57
3.2	Physical parameters of cultivation of <i>B. sp.</i> under natural conditions	57
3.3	Reagent Composition for Gel.	63
4.1	Elemental composition of <i>Botryococcus</i> sp	73
4.2	Proximate analysis of <i>B. sp.</i> biomass under nitrogen atmospheres. Values are reported as the mean \pm standard deviation	74
4.3	The result from the ultimate analysis of <i>Botryococcus</i> sp biomass. Values are reported as the mean \pm standard deviation	75
4.4	Composition of monomeric sugars in <i>Botryococcus</i> sp	77
4.5	Fatty acids composition of <i>Botryococcus</i> sp.	82
4.6	Fatty acid profile and content (%) of some microalgae	83
4.7	Extracellular metabolites of <i>B. sp.</i> obtained from the BBM	86
4.8	Identify enzymes related to carbohydrates metabolism	92
4.9	The blastp result shows that it is Query_316718	94

LIST OF FIGURES

2.1	Photosynthesis mechanisms of microalgae influenced by light intensity, photoperiod, temperature, and salinity	15
2.2	Calvin cycle :Adapted from (Bari, 2004)	17
2.3	Carbohydrate metabolism of green algae Adapted from (Chen <i>et al.</i> , 2013)	38
2.4	Simplified overview of the metabolites and representative pathways in microalgal lipid biosynthesis shown in black and enzymes demonstrated in red Adapted from (Radakovits <i>et al.</i> , 2010)	42
3.1	Casting frames clamped to casting stand	63
3.2	Samples loaded into individual wells.	64
3.3	Electrophoresis of the protein.	65
3.4	Gel staining in Coomassie blue	65
3.5	Gel inside the destaining solution.	66
4.1	The growth rate of <i>B. sp.</i> in the bald basal medium (BBM)	70
4.2	Light microscopic image of <i>B. sp</i> ($\times 40$) observed under a light compound microscope.	71
4.3	SEM images ($5000\times$) of <i>B. sp.</i> cells.	71
4.4	EDX spectrum of <i>Botryococcus sp.</i>	73
4.5	TG profile of <i>Botryococcus sp.</i>	76
4.6	DTG profile of <i>Botryococcus sp.</i>	76
4.7	HPLC Chromatogram of <i>B. sp.</i> Sugars	78
4.8	Monomeric sugars in <i>Botryococcus sp</i>	79
4.9	SDS-PAGE analysis of the major protein fractions from <i>B. sp.</i> (arrows showing apparent bands).	81
4.10	GC-MS chromatogram of <i>B. sp.</i> Supernatant	85

4.11	Graphical illustration of the Constructed alternative pathway of <i>Botryococcus</i> sp.	91
------	---	----



LIST OF ABBREVIATIONS

ATP	- Adenosine Triphosphate
APS	- Ammonium Persulfate
BBC	- Bold Basal Medium
BSA	- Bovine Serum Albumin
CC	- Calvin Cycle
CO ₂	- Carbon IV Oxide
CO ₃ ²⁻	- Carbonate
CaCl ₂	- Calcium Chloride
DDT	- Dithiothreitol
DDH ₂ O	- Double Distilled Water
DHA	- Docosahearnoi Acid
DNA	- Dioxyribonucleic Acid
EDX	- Energy Dispensive X- Ray
ER	- Endoplasmic Reticulum
EPA	- Eicosapentaenoicacid
ESP	- ExoPolysaccharide
GHG	- Green House Gases
H ₂ PO ₄ ⁻	- hydrogen phosphate ion
HPO ₄ ²⁻	- hydrogen phosphate
HPLC	- High Performance Liquid Chromatography
IAA	- Iodoaetamide
LC- MS	- Liquid Chromatography Mass Spectrometer
MG	- Magnesium
NaCl	- Sodium Chloride
NADP ⁺	- Nicotinamide Adenine dinucleotide Phosphate
NaNO ₃	- Sodium Nitrate
NREL	- Renewable Energy Laboratory

NO ₃	- Nitrate
O ₂	- Oxygen
P	- Phosphorus
PRB	- Photo Bioreactor
PGA	- Phosphoric Acid
PUFA	- Polyunsaturated Fatty Acid
PAGE	- Polyacrylamide Gel Electrophoresis
RUBISCO	- Ribulose 1, 5 - Biphosphate carboxylase
RNA	- Ribonucleic Acid
SI	- Siloxane
SDS	- Sodium Chloride Sulphate
TAG	- Triacygly Ceride or Triglycerides
TEMED	- Tetrametylethylenediamine
UTHM	- Universiti Tun Hussein Onn Malaysia

-



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

LIST OF APPENDICIES

APPENDICIES	TITLE	PAGE
A	Botryococcus cultivation	131
B	Specific growth rate	132
C	The result of molecular analysis	133
D	The migration distance of protein bands	136
E	GC- MS chromatograms	138
F	Enzymes	139
G	Carbohydrate monomers	153
H	BLAST Result	156
I	List of publication	161



CHAPTER 1

INTRODUCTION

1.1 Background

Currently, fossil fuels provide about 80 % of global energy demand (Petroleum, 2012). Issues such as decreasing reserves, inadequate supply, and unavoidably growing market, the world's cheapest source of energy is still fossil fuels. However, the prices of fossil fuels have increased as demand has exceeded supply. Therefore, many nations have started to ponder the problem of energy safety, mostly as fossil fuels are their only energy source. Consequently, widespread exploitation of fossil fuels has caused organic greenhouse gases (GHG) release, in which more than 60% is carbon dioxide that is mostly responsible for global climate change. Furthermore, the environmental impact has an enormous effect on the carbon cycle (carbon balance), which is related to the combustion of fossil fuels. Besides, exhaustion of different existing biomass without appropriate compensation resulted in colossal biomass scarcity, developing environmental problems like deforestation and the destruction of biodiversity.

Improving the rate of reduction of fossil fuels and reducing GHG emissions needs sufficient renewable clean energy to be generated for use. Small to medium-term renewable and environmentally friendly alternatives for fossil fuels have lately attracted the serious attention of sciences worldwide (Ghaffour *et al.*, 2015).

Among several possible options, at present is biofuels. This is one of the utmost interest and are likely to show a significant role in global energy substructure. Biofuels are fuels that comprise energy from geologically current carbon fixation by living organisms. Biofuels produced from biomasses are usually from plants or materials originating from plants or biomasses from algae and microalgae, which are

biodegradable, non-toxic and renewable (Ioelovich, 2013, 2015; Song *et al.*, 2008). They use solar energy, together with carbon dioxide, to convert and store the biomass in the forms of carbohydrates, lipids, and proteins through the process of photosynthesis (Liu *et al.*, 2015).

Microalgae is a unicellular organism that executes higher photosynthetic effectiveness and growth potential compared with terrestrial plants (Katiyar *et al.*, 2017). Since it has high photosynthetic efficiency, the mass cultivation can reduce the carbon dioxide emission efficiently from the atmosphere, hence, decreasing the effect of global warming (Surriya *et al.*, 2015). According to Milano *et al.* (2016), microalgae have a high growth rate and can produce up to 70 % of lipid content inside their cells depend on the type of species. They also reported that microalgae can survive under the severe condition and can grow in a small cultivation land area. Its cultivation does not compete with land plants with the growth of crops for food and biofuel and never require rigorous treatment methods in comparison with lignocellulose-enriched biomass. Furthermore, microalgae can grow on nonagricultural soil using wastewater instead of drinking water (Gani *et al.*, 2016). They accumulate carbohydrates and lipids, which are the most abundant constituents of microalgal cells, in addition to other valuable chemicals, such as carotenoids and proteins (Jeon *et al.*, 2017).

The existence of the microalga *Botryococcus* sp. in America, Africa, Asia and Europe has confirmed its wide distribution (Chandra, 1964; Chandrappa *et al.*, 2010; Metzger *et al.*, 1985; Okada *et al.*, 2000; Wolf *et al.*, 1985). Furthermore, these geographical regions belong to different climatic zones like continental, temperate, tropical and alpine, hence an indication of its ability to grow in varied climatic conditions (Tyson, 2012). The ability of microalgae to survive in diverse and extreme conditions is reflected in its tremendous diversity and also the unusual pattern of cellular lipids obtained from these microalgae (Sato *et al.*, 2000). Furthermore, some of these microalgae can also modify lipid metabolism efficiently to counter adverse changes in environmental conditions (Guschina & Harwood, 2006; Thompson Jr, 1996). The most prominent characteristic of *Botryococcus* is its ability to synthesize and accumulate very high levels of lipids. These lipid substances include numerous hydro-carbons, that is, highly reduced compounds comprising of only carbon and hydrogen as elements (Brown *et al.*, 1969). *Botryococcus* being a photosynthetic organism, has been reported to reduce CO₂ emissions by 1.5×10^5 tons/yr./ 8.4×10^3

ha (Sawayama *et al.*, 1999) thereby offering an eco-friendly process for production of lipid and other bioactive compounds along with its carbon dioxide mitigation credits.

Additionally, *Botryococcus* spp. are well known to have a remarkable capability to synthesise industrially interesting chemicals intracellular and also excrete them into the extracellular matrix. Although *Chlorella* and *Dunaliella* are both highly efficient microalgae with regards to growth-to-lipid ratio, *Botryococcus* has outclassed them both. *Botryococcus* can produce a large number of hydrocarbons as compared to its biomass (Chisti, 2007a; Cohen, 2014). The production level of oil content in microalga *Chlorella* is up to 50% lipids, while *Botryococcus* produces the highest oil content of approximately 80% (Powell & Hill, 2009).

Due to its potential to produce large amounts of lipids and hydrocarbons, exploration of the genus *Botryococcus* is still ongoing for newer strains and species and to date, more than 60 *Botryococcus* strains have been cultivated in laboratories (For a non-exhaustive list, see Metzger and Largeau, 1999) and the exploration is still on. Most *Botryococcus* strains cultivated in laboratories and wild samples collected from lakes worldwide are analyzed for their lipid production (Metzger and Largeau, 1999) and are reported to produce 25 - 75% of lipids on their dry weight (Metzger and Largeau, 1999, 2005; Dayananda *et al.*, 2007a; Ranga Rao *et al.*, 2007; Hai-Linh *et al.*, 2009; Ela and Anastasios, 2010; Chiara *et al.*, 2010).

However, there are no reports available regarding why and what is responsible for the high lipid content in *B. sp.* in comparison to other microalgae. In view of its importance, the present study aims to find out the cause from an indigenous species of *Botryococcus* isolated in TJNER.

To enhance the economic feasibility of mass production of micro-algal biofuels, several efforts are still being made to increase strain performance (Xin *et al.*, 2010), cultivation methods (Ketheesan & Nirmalakhandan, 2012; Zhang & Hu, 2012) and improving technologies involved in biomass harvesting and extraction (Halim *et al.*, 2012; Hanotu *et al.*, 2012).

1.2 Statement of the problem

The rising interest surrounding the continued use of fossil fuels and rapid reduction of fossil fuel reserves, global climate change, increasing crude oil price and

environmental degradation has enforced governments, policymakers, scientists, and researchers worldwide to find alternative renewable energy sources like biofuels (Alam *et al.*, 2015; Dragone *et al.*, 2010). Microalgae have emerged as a promising feedstock for the production of biofuels leading to explorations for a suitable microalgal strain for biotechnological exploitation with a higher lipid content than what is obtainable now.

Botryococcus spp. have been reported to be richer in lipid content (Huang *et al.* 2013) than other microalgae with also a significant high lipid content for example, *Chlorella vulgaris* and *Dunaliella* (Griffiths and Harrison 2009; Keffer and Kleinheinz 2002). This begs the question of why or what is the cause of this disparity in lipid content between *Botryococcus* and other microalgal species.

Furthermore, according to Cakmak *et al.* (2012), Hu *et al.* (2008), Li *et al.* (2011) and Siaut *et al.* (2011), photosynthetic carbon partitioning switch towards the production of energy-rich storage compounds like starch and lipid, in which it can be converted into biofuels is possible. In other words, the carbohydrate present in cells can be converted to lipids by manipulating the carbohydrate pathway to increase the lipid content, hence maximising the overall biofuel production.

Therefore, understanding the metabolic pathways and processes involved in the generation of these organic macromolecules is essential to improve biofuels production. However, studies on carbohydrate metabolic pathway and enzymes discovery leading to the production of lipid in *B. sp.* are minimal. Hence the need to study the metabolic pathways and identify the enzymes involved in the metabolism of carbohydrates in *B. sp.* that will lead to more lipid production with the hope of improving its biofuel productivity. Conversely, microalgae have a strategic feature of rapid alteration of the intracellular energy storage form from starch to lipids, thereby generating more lipids contents. Overall, this is a fundamental study into the basic composition and metabolic biosynthesis of the microalga *Botryococcus* sp.

1.3 Research aim and objectives

This research work aims to determine the metabolic pathway and the enzymes involved in carbohydrate metabolism of *B. sp.* The objectives of this study include:

- ❖ To identify the intercellular and extracellular metabolites produce from *B. sp.*

cultured under natural conditions.

- ❖ To evaluate the enzymes involved in carbohydrate metabolism leading to lipid production in *B. sp.* under natural conditions.
- ❖ To construct an alternative carbohydrate to lipid metabolic pathway of *B. sp.* cultivated under natural conditions.

1.4 The importance of the study

Renewable and maintainable energy sources from photosynthetic microalgae have received much attention in recent years. *B. sp.* is capable of converting sunlight and CO₂ into biomass and other valuable products. *B. sp.* was initially collected from Endau Rompin National Park and known as a wild type of microalga. The microalgae are an indigenous species and have not been studied intensively yet. Because of its virtually unique capability to synthesise large quantities of exopolysaccharides and hydrocarbons, this colony-forming green microalga serves as a potential source for industrial applications like biofuels and other valuable chemicals.

However, there is little understanding of the metabolism that will channel polysaccharides to this lipid production. Through studying the intracellular and extracellular metabolites, it enabled us to know the mass balance inside and outside of the microalgae since we understand what it consumes. This study also contributes to giving much understanding on the mechanism of the carbohydrate pathway based on the identified proteins/enzymes as a primary novelty to enhance the potential of biofuel production in future.

This study will also aim to establish an alternative pathway of lipid synthesis which will not be based on the genetic modification but the export of the fixed carbon (3PGA) to the cytoplasm, providing a shorter route to lipid production than the normal process via the plastid. This pathway will prove to be a different route to producing more lipid that was not established before. The alternate pathway will also show the main reason why the *B. sp.* was able to produce more lipids than carbohydrate.

1.5 The scope of the study

This research was carried out to determine and identify the enzymes involved in carbohydrate metabolism leading to lipid production and to construct this pathway subsequently. The study covered the culture *B. sp.* isolated from the rocky pools of Taman Negara Endau Rompin, Johor, Malaysia. The microalgae were mass cultured in Bold basal medium (BBM) at an ambient temperature of 25 °C to 30 °C pending on the time of the day within the lumination of 30-60 W/M² from sunlight, harvested and freeze-dried. Proximate analysis to determine the percentage of carbohydrates, lipid and proteins contents was carried out. Additionally, the monomeric composition of sugars, fatty acids, characterisation of extracted proteins to identify enzymes and the metabolites produced by the microalga were determined. Finally, the carbohydrate pathway was constructed based on the information gathered from the previous analysis.



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

CHAPTER 2

LITERATURE REVIEW

2.1 Microalgae, *Botryococcus* sp.

Microalgae are considered to be the oldest living organisms on the planet (Groombridge *et al.*, 2002; Richmond, 2008). They are the primary producers and can fix nearly 50 % of the global carbon (77 %) with the production of oxygen, O₂ (Metz *et al.*, 2007). Rawat *et al.* (2013) reported that more than 50,000 microalgae species existing in the world but estimated only 30,000 identified. Depending on the species of microalgae, their sizes ranged from micrometres to millimetres. The term microalgae have no formal taxonomic standing and are defined as thallophytes lacking roots, stems, and leaves with chlorophyll as their primary photosynthetic pigment (Brennan & Owende, 2010; Kumar *et al.*, 2013; Saifullah *et al.*, 2014). They are prokaryotic organisms with simple cellular structure and a large surface to volume body ratio, which offers them the capacity to uptake a lot of nutrients (Fogg, 2012). They grow very fast and live in different environmental conditions such as the aquatic and terrestrial ecosystems (Mata *et al.*, 2010; Saifullah *et al.*, 2014).

The green algae are one of the largest algae groups regarding species number, biological distribution, and morphological diversity. The eukaryotic algal species, the green algae are found in an enormous variety of habitats (Wehr *et al.*, 2015). They occur in lakes, freshwater, streams, and ponds also on and in the soil, rocks, snow, ice, plants, and animals. Lastly, algae can also be free-living or be in a relationship with other organisms, such as lichens (Sankaran & Thiruneelagandan, 2015). Morphological differentiation is ranging from unicellular strains (free-living

REFERENCES

- Abdel-Raouf, N., Al-Homaidan, A., & Ibraheem, I. (2012). Microalgae and wastewater treatment. *Saudi journal of biological sciences*, 19(3), 257-275.
- Adrio, J. L., & Demain, A. L. (2014). Microbial enzymes: tools for biotechnological processes. *Biomolecules*, 4(1), 117-139.
- Alabi, A., Tampier, M., & Bibeau, E. (2009). Microalgae technologies for biofuels/bioenergy production in British Columbia: Current technology and barriers to implementation: Final report to the British Columbia Innovation Council. *Seed Science Ltd. Report (Seed Sci., 2009)*. http://www.fao.org/uploads/media/0901_Seed_Science_Microalgae_technologies_and_processes_for_biofuelsbioenergy_production_in_British_Columbia.pdf. Google Scholar.
- Alam, F., Mobin, S., & Chowdhury, H. (2015). Third generation biofuel from Algae. *Procedia Engineering*, 105, 763-768.
- Allard, B., & Casadevall, E. (1990). Carbohydrate composition and characterization of sugars from the green microalga *Botryococcus braunii*. *Phytochemistry*, 29(6), 1875-1878.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of molecular biology*, 215(3), 403-410.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research*, 25(17), 3389-3402.
- Andersen, R. A. (2005). *Algal culturing techniques*: Elsevier.
- Apha, A. (2012). WEF.(2012). *Standard methods for the examination of water and wastewater*, 22.
- Apt, K. E., & Behrens, P. W. (1999). Commercial developments in microalgal biotechnology. *Journal of phycology*, 35(2), 215-226.

- Arguelles, E. D., Laurena, A. C., Monsalud, R. G., & Martinez-Goss, M. R. (2018). Fatty acid profile and fuel-derived physico-chemical properties of biodiesel obtained from an indigenous green microalga, *Desmodesmus sp.(I-AU1)*, as potential source of renewable lipid and high quality biodiesel. *Journal of Applied Phycology*, 30(1), 411-419.
- Armbrust, E. V. (1999). Identification of a New Gene Family Expressed during the Onset of Sexual Reproduction in the Centric Diatom *Thalassiosira weissflogii*. *Applied and environmental microbiology*, 65(7), 3121-3128.
- Ashokkumar, V., & Rengasamy, R. (2012). Mass culture of *Botryococcus braunii* Kutz. under open raceway pond for biofuel production. *Bioresource technology*, 104, 394-399.
- Avinash, A., Subramaniam, D., & Murugesan, A. (2014). Bio-diesel—A global scenario. *Renewable and Sustainable Energy Reviews*, 29, 517-527.
- Badawy, M., Abou-Waly, H., & Ali, G. (1999). Excretion products of algae and their occurrence in Solar Lake. Taba, Egypt. *International Journal of Environmental Health Research*, 9(3), 233-243.
- Bafana, A. (2013). Characterization and optimization of production of exopolysaccharide from *Chlamydomonas reinhardtii*. *Carbohydrate polymers*, 95(2), 746-752.
- Bai, F., Anderson, W., & Moo-Young, M. (2008). Ethanol fermentation technologies from sugar and starch feedstocks. *Biotechnology advances*, 26(1), 89-105.
- Banerjee, A., Sharma, R., Chisti, Y., & Banerjee, U. (2002). *Botryococcus braunii*: a renewable source of hydrocarbons and other chemicals. *Critical reviews in biotechnology*, 22(3), 245-279.
- Barbarino, E., & Lourenço, S. O. (2005). An evaluation of methods for extraction and quantification of protein from marine macro-and microalgae. *Journal of Applied Phycology*, 17(5), 447-460.
- Barbosa, M. J., & Wijffels, R. H. (2004). Overcoming shear stress of microalgae cultures in sparged photobioreactors. *Biotechnol Bioeng*, 85(1), 78-85.
- Bari, R. (2004). *A novel approach for the suppression of photorespiration in C3 plants by gene transfer*. Bibliothek der RWTH Aachen.
- Baroukh, C., Muñoz-Tamayo, R., Steyer, J.-P., & Bernard, O. (2015). A state of the art of metabolic networks of unicellular microalgae and cyanobacteria for biofuel production. *Metabolic engineering*, 30, 49-60.

- Barsanti, L., & Gualtieri, P. (2014). *Algae: anatomy, biochemistry, and biotechnology*: CRC press.
- Bauwe, H., Hagemann, M., Kern, R., & Timm, S. (2012). Photorespiration has a dual origin and manifold links to central metabolism. *Current opinion in plant biology*, 15(3), 269-275.
- Bayona, K. C. D., & Garcés, L. A. (2014). Effect of different media on exopolysaccharide and biomass production by the green microalga *Botryococcus braunii*. *Journal of Applied Phycology*, 26(5), 2087-2095.
- Becker, E. (2007). Micro-algae as a source of protein. *Biotechnology advances*, 25(2), 207-210.
- Becker, E. W. (1994). *Microalgae: biotechnology and microbiology* (Vol. 10): Cambridge University Press.
- Belcher, H., & Swale, E. (1976). *A beginner's guide to freshwater algae*: HMSO.
- Berg, A. H., Combs, T. P., & Scherer, P. E. (2002). ACRP30/adiponectin: an adipokine regulating glucose and lipid metabolism. *Trends in Endocrinology & Metabolism*, 13(2), 84-89.
- Berman-Frank, I., & Dubinsky, Z. (1999). Balanced growth in aquatic plants: myth or reality? Phytoplankton use the imbalance between carbon assimilation and biomass production to their strategic advantage. *Bioscience*, 49(1), 29-37.
- Bhat, M., & Bhat, S. (1997). Cellulose degrading enzymes and their potential industrial applications. *Biotechnology advances*, 15(3-4), 583-620.
- Bi, Z., & He, B. B. (2013). Characterization of microalgae for the purpose of biofuel production. *Transactions of the ASABE*, 56(4), 1529-1539.
- Bišová, K., & Zachleder, V. (2014). Cell-cycle regulation in green algae dividing by multiple fission. *Journal of experimental botany*, 65(10), 2585-2602.
- Blifernez-Klassen, O., Chaudhari, S., Klassen, V., Wördenweber, R., Steffens, T., Cholewa, D., Kruse, O. (2018). Metabolic survey of *Botryococcus braunii*: Impact of the physiological state on product formation. *PloS one*, 13(6), e0198976.
- Borowitzka, M. A. (1995). Microalgae as sources of pharmaceuticals and other biologically active compounds. *Journal of Applied Phycology*, 7(1), 3-15.
- Borowitzka, M. A. (2013). Energy from microalgae: a short history *Algae for biofuels and energy* (pp. 1-15): Springer.

- Borowitzka, M. A., Beardall, J., & Raven, J. A. (2016). *The physiology of microalgae* (Vol. 6): Springer.
- Bouterfas, R., Belkoura, M., & Dauta, A. (2002). Light and temperature effects on the growth rate of three freshwater [2pt] algae isolated from a eutrophic lake. *Hydrobiologia*, 489(1-3), 207-217.
- Bouterfas, R., Belkoura, M., & Dauta, A. (2006). The effects of irradiance and photoperiod on the growth rate of three freshwater green algae isolated from a eutrophic lake. *Limnetica*, 25(3), 647-656.
- Boyd, C. (1973). Amino acid composition of freshwater algae. *Arch Hydrobiol*, 72, 1-9.
- Brányiková, I., Maršáľková, B., Doucha, J., Brányik, T., Bišová, K., Zachleder, V., & Vítová, M. (2011). Microalgae—novel highly efficient starch producers. *Biotechnol Bioeng*, 108(4), 766-776.
- Brasil, B. d. S. A. F., de Siqueira, F. G., Salum, T. F. C., Zanette, C. M., & Spier, M. R. (2017). Microalgae and cyanobacteria as enzyme biofactories. *Algal Research*, 25, 76-89.
- Brennan, L., & Owende, P. (2010). Biofuels from microalgae—a review of technologies for production, processing, and extractions of biofuels and co-products. *Renewable and Sustainable Energy Reviews*, 14(2), 557-577.
- Breuer, G., de Jaeger, L., Artus, V. G., Martens, D. E., Springer, J., Draaisma, R. B., . . . Lamers, P. P. (2014). Superior triacylglycerol (TAG) accumulation in starchless mutants of *Scenedesmus obliquus*:(II) evaluation of TAG yield and productivity in controlled photobioreactors. *Biotechnology for biofuels*, 7(1), 70.
- Breuer, G., Lamers, P. P., Martens, D. E., Draaisma, R. B., & Wijffels, R. H. (2013). Effect of light intensity, pH, and temperature on triacylglycerol (TAG) accumulation induced by nitrogen starvation in *Scenedesmus obliquus*. *Bioresource technology*, 143, 1-9.
- Brown, A., Knights, B., & Conway, E. (1969). Hydrocarbon content and its relationship to physiological state in the green alga *Botryococcus braunii*. *Phytochemistry*, 8(3), 543-547.
- Carneiro, C. R., Postol, E., Nomizo, R., Reis, L. F., & Brentani, R. R. (2004). Identification of enolase as a laminin-binding protein on the surface of *Staphylococcus aureus*. *Microbes and Infection*, 6(6), 604-608.

- Carriquiry, M. A., Du, X., & Timilsina, G. R. (2011). Second generation biofuels: Economics and policies. *Energy Policy*, 39(7), 4222-4234.
- Casadevall, E., Dif, D., Largeau, C., Gudin, C., Chaumont, D., & Desanti, O. (1985). Studies on batch and continuous cultures of *Botryococcus braunii*: hydrocarbon production in relation to physiological state, cell ultrastructure, and phosphate nutrition. *Biotechnol Bioeng*, 27(3), 286-295.
- Castillo-Romero, A., Davids, B. J., Lauwaet, T., & Gillin, F. D. (2012). Importance of enolase in *Giardia lamblia* differentiation. *Molecular and biochemical parasitology*, 184(2), 122-125.
- Chaisutyakorn, P., Praiboon, J., & Kaewsuralikhit, C. (2018). The effect of temperature on growth and lipid and fatty acid composition on marine microalgae used for biodiesel production. *Journal of Applied Phycology*, 30(1), 37-45.
- Chandra, A. (1964). A note on the occurrence of *Botryococcus* in the miocene lignites of Kerala. *Current Science*, 33(7), 214-215.
- Chandrappa, D., Lakshmanan, V., Neelwarne, B., & Aswathanarayana, R. G. (2010). Assessment of genetic polymorphism among green microalgae *Botryococcus* of distinct origin by RAPD. *Genes Genomes and Genomics*, 4, 65-69.
- Chang, R. L., Ghamsari, L., Manichaikul, A., Hom, E. F., Balaji, S., Fu, W., Salehi-Ashtiani, K. (2011). Metabolic network reconstruction of *Chlamydomonas* offers insight into light-driven algal metabolism. *Molecular systems biology*, 7(1), 518.
- Chatsungnoen, T., & Chisti, Y. (2016). Oil production by six microalgae: impact of flocculants and drying on oil recovery from the biomass. *Journal of Applied Phycology*, 28(5), 2697-2705.
- Checchetto, V., Teardo, E., Carraretto, L., Formentin, E., Bergantino, E., Giacometti, G. M., & Szabo, I. (2013). Regulation of photosynthesis by ion channels in cyanobacteria and higher plants. *Biophysical chemistry*, 182, 51-57.
- Chen, C.-Y., Zhao, X.-Q., Yen, H.-W., Ho, S.-H., Cheng, C.-L., Lee, D.-J., Chang, J.-S. (2013). Microalgae-based carbohydrates for biofuel production. *Biochemical Engineering Journal*, 78, 1-10.
- Chen, C. Y., & Durbin, E. G. (1994). Effects of pH on the growth and carbon uptake of marine phytoplankton. *Marine Ecology Progress Series*, 83-94.

- Chen, M., Chen, X., Liu, T., & Zhang, W. (2011). Subcritical ethanol extraction of lipid from wet microalgae paste of *Nannochloropsis* sp. *Journal of Biobased Materials and Bioenergy*, 5(3), 385-389.
- Cheng, D., & He, Q. (2014). Assessment of environmental stresses for enhanced microalgal biofuel production—an overview. *Frontiers in Energy Research*, 2, 26.
- Cheng, P., Zhou, C., Wang, Y., Xu, Z., Xu, J., Zhou, D., Liu, T. (2018). Comparative transcriptome analyses of oleaginous *Botryococcus braunii* race A reveal significant differences in gene expression upon cobalt enrichment. *Biotechnology for biofuels*, 11(1), 333.
- Cheng, Y. S., Labavitch, J., & VanderGheynst, J. (2015). Elevated CO₂ concentration impacts cell wall polysaccharide composition of green microalgae of the genus *Chlorella*. *Letters in applied microbiology*, 60(1), 1-7.
- Chepurnov, V. A., Mann, D. G., Von Dassow, P., Vanormelingen, P., Gillard, J., Inze, D., Vyverman, W. (2008). In search of new tractable diatoms for experimental biology. *BioEssays*, 30(7), 692-702.
- Chiranjeevi, P., & Mohan, S. V. (2016). Critical parametric influence on microalgae cultivation towards maximizing biomass growth with simultaneous lipid productivity. *Renewable energy*, 98, 64-71.
- Chisti, Y. (2007a). Biodiesel from microalgae. *Biotechnology advances*, 25(3), 294-306.
- Chisti, Y. (2007b). Biodiesel from microalgae *Biotechnology Advances* 25: 294–306. *Google Scholar*.
- Cohen, Z. (2014). *Chemicals from microalgae*: CRC Press.
- Contreras, L., Ritter, A., Dennett, G., Boehmwald, F., Guitton, N., Pineau, C., Correa, J. A. (2008). Two-dimensional gel electrophoresis analysis of brown algal protein extracts 1. *Journal of phycology*, 44 (5), 1315-1321.
- Da-Cong, Y., Ya-Hong, G., Hong, M., Zheng-Rong, O., Hong-Jun, H., & Ye-Guang, L. (2008). The Effects of Several Environmental Factors on the Photosynthesis of *Botryococcus braunii*. *Journal of Wuhan Botanical Research*, 1, 012.
- Danquah, M. K., Gladman, B., Moheimani, N., & Forde, G. M. (2009). Microalgal growth characteristics and subsequent influence on dewatering efficiency. *Chemical Engineering Journal*, 151 (1-3), 73-78.

- Davies, C. M., Apte, S. C., Peterson, S. M., & Stauber, J. L. (1994). Plant and algal interference in bacterial beta-D-galactosidase and beta-D-glucuronidase assays. *Applied and environmental microbiology*, 60 (11), 3959-3964.
- Dayan, C., Kumudha, A., Sarada, R., & Ravishankar, G. (2010). Isolation, characterization and outdoor cultivation of green microalgae *Botryococcus* sp. *Scientific Research and Essays*, 5 (17), 2497-2505.
- Dayananda, C., Sarada, R., Rani, M. U., Shamala, T., & Ravishankar, G. (2007). Autotrophic cultivation of *Botryococcus braunii* for the production of hydrocarbons and exopolysaccharides in various media. *Biomass and Bioenergy*, 31(1), 87-93.
- De Moraes, M. G., & Costa, J. A. V. (2007). Biofixation of carbon dioxide by *Spirulina* sp. and *Scenedesmus obliquus* cultivated in a three-stage serial tubular photobioreactor. *Journal of biotechnology*, 129(3), 439-445.
- de Moraes, M. G., Vaz, B. d. S., de Moraes, E. G., & Costa, J. A. V. (2015). Biologically active metabolites synthesized by microalgae. *BioMed research international*, 2015.
- de Oliveira Dal'Molin, C. G., Quek, L.-E., Palfreyman, R. W., & Nielsen, L. K. (2011). *AlgaGEM—a genome-scale metabolic reconstruction of algae based on the Chlamydomonas reinhardtii genome*. Paper presented at the BMC genomics.
- de Oliveira Rangel-Yagui, C., Danesi, E. D. G., de Carvalho, J. C. M., & Sato, S. (2004). Chlorophyll production from *Spirulina platensis*: cultivation with urea addition by fed-batch process. *Bioresource technology*, 92 (2), 133-141.
- de Siqueira, F. G., Ferreira, F., & Ximenes, E. (2010). Plant cell wall as a substrate for the production of enzymes with industrial applications. *Mini-Reviews in Organic Chemistry*, 7(1), 54-60.
- de Souza, M. F., Pereira, D. S., Freitas, S. P., da Silva Bon, E. P., & Rodrigues, M. A. (2017). Neutral sugars determination in *Chlorella*: Use of a one-step dilute sulfuric acid hydrolysis with reduced sample size followed by HPAEC analysis. *Algal Research*, 24, 130-137.
- Delbarre-Ladrat, C., Sinquin, C., Lebellenger, L., Zykwinska, A., & Collic-Jouault, S. (2014). Exopolysaccharides produced by marine bacteria and their applications as glycosaminoglycan-like molecules. *Frontiers in chemistry*, 2, 85.

- Dertli, E., Mayer, M. J., & Narbad, A. (2015). Impact of the exopolysaccharide layer on biofilms, adhesion and resistance to stress in *Lactobacillus johnsonii* FI9785. *BMC microbiology*, 15(1), 8.
- Deschamps, P., Colleoni, C., Nakamura, Y., Suzuki, E., Putaux, J.-L., Buléon, A., Falcón, L. I. (2007). Metabolic symbiosis and the birth of the plant kingdom. *Molecular Biology and Evolution*, 25(3), 536-548.
- Deschamps, P., Haferkamp, I., Dauvillée, D., Haebel, S., Steup, M., Buléon, A., lancke, C. (2006). Nature of the periplastidial pathway of starch synthesis in the cryptophyte *Guillardia theta*. *Eukaryotic cell*, 5(6), 954-963.
- Dey, P., & Kauss, H. (1981). α -galactosidase of *Poterioochromonas malhamensis*. *Phytochemistry*, 20(1), 45-48.
- Díaz-Ramos, À., Roig-Borrellas, A., García-Melero, A., & López-Aleman, R. (2012). α -Enolase, a multifunctional protein: its role on pathophysiological situations. *BioMed research international*, 2012.
- Ding, J., Zhao, F., Cao, Y., Xing, L., Liu, W., Mei, S., & Li, S. (2015). Cultivation of microalgae in dairy farm wastewater without sterilization. *International journal of phytoremediation*, 17(3), 222-227.
- Dragone, G., Fernandes, B. D., Vicente, A. A., & Teixeira, J. A. (2010). Third generation biofuels from microalgae. *Current research, technology and education topics in applied microbiology and microbial biotechnology*, 2, 1355-1366.
- Du, Z., Hu, B., Ma, X., Cheng, Y., Liu, Y., Lin, X., Ruan, R. (2013). Catalytic pyrolysis of microalgae and their three major components: carbohydrates, proteins, and lipids. *Bioresource technology*, 130, 777-782.
- Dudoladova, M. V., Kupriyanova, E. V., Markelova, A. G., Sinetova, M. P., Allakhverdiev, S. I., & Pronina, N. A. (2007). The thylakoid carbonic anhydrase associated with photosystem II is the component of inorganic carbon accumulating system in cells of halo-and alkaliphilic cyanobacterium *Rhabdoderma lineare*. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1767 (6), 616-623.
- Elzenga, J. T. M., Prins, H., & Stefels, J. (2000). The role of extracellular carbonic anhydrase activity in inorganic carbon utilization of *Phaeocystis globosa* (Prymnesiophyceae): a comparison with other marine algae using the isotopic disequilibrium technique. *Limnology and Oceanography*, 45 (2), 372-380.

- Ester, D., & Pramoda, J. (2013). A Short Review on Proteomics and its Applications. *International Letters of Natural Sciences*.
- Fagiri, Y. M. A., Salleh, A., & El-Nagerabi, S. A. (2013). Influence of chemical and environmental factors on the growth performance of *Spirulina platensis* strain SZ100. *J Algal Biomass Utiln*, 4(2), 7-15.
- Feist, P., & Hummon, A. B. (2015). Proteomic challenges: sample preparation techniques for microgram-quantity protein analysis from biological samples. *International journal of molecular sciences*, 16(2), 3537-3563.
- Feldmane, J., Semjonovs, P., & Ciprovica, I. (2013). *Potential of exopolysaccharides in yoghurt production*. Paper presented at the Proceedings of World Academy of Science, Engineering and Technology.
- Fernandes, B., Teixeira, J., Dragone, G., Vicente, A. A., Kawano, S., Bišová, K., . . . Vítová, M. (2013). Relationship between starch and lipid accumulation induced by nutrient depletion and replenishment in the microalga *Parachlorella kessleri*. *Bioresource technology*, 144, 268-274.
- Fernandes, H., Tomé, M., Lupi, F., Fialho, A., Sá-Correia, I., & Novais, J. (1989). Biosynthesis of high concentrations of an exopolysaccharide during the cultivation of the microalga *Botryococcus braunii*. *Biotechnology Letters*, 11(6), 433-436.
- Fogg, G. (2012). *The blue-green algae*: Elsevier.
- Ganapathy, K., Chidambaram, K., Janarthanan, R., & Ramasamy, R. (2017). Effect of UV-B radiation on growth, photosynthetic activity and metabolic activities of *Chlorella vulgaris*. *Journal of Microbiology and Biotechnology*, 6, 53-60.
- Gani, P., Mohamed Sunar, N., Matias-Peralta, H., Abdul Latiff, A. A., & Mohamad Fuzi, S. F. Z. (2017a). Growth of microalgae *Botryococcus* sp. in domestic wastewater and application of statistical analysis for the optimization of flocculation using alum and chitosan. *Preparative Biochemistry and Biotechnology*, 47(4), 333-341.
- Gani, P., Sunar, N., Matias-Peralta, H., Latiff, A. A., & Razak, A. (2016). Influence of initial cell concentrations on the growth rate and biomass productivity of microalgae in domestic wastewater. *Appl. Ecol. Environ. Res*, 14(2), 399-409.
- Gani, P., Sunar, N. M., Matias-Peralta, H., Mohamed, R. M. S. R., Latiff, A. A. A., & Parjo, U. K. (2017b). Extraction of hydrocarbons from freshwater green

- microalgae (*Botryococcus* sp.) biomass after phycoremediation of domestic wastewater. *International journal of phytoremediation*, 19(7), 679-685.
- Gardner, R. D., Lohman, E., Gerlach, R., Cooksey, K. E., & Peyton, B. M. (2013). Comparison of CO₂ and bicarbonate as inorganic carbon sources for triacylglycerol and starch accumulation in *Chlamydomonas reinhardtii*. *Biotechnol Bioeng*, 110(1), 87-96.
- Garibay-Hernández, A., Barkla, B. J., Vera-Estrella, R., Martinez, A., & Pantoja, O. (2017). Membrane proteomic insights into the physiology and taxonomy of an oleaginous green microalga. *Plant physiology*, 173(1), 390-416.
- Gavin, A.-C., Bösch, M., Krause, R., Grandi, P., Marzioch, M., Bauer, A., Cruciat, C.-M. (2002). Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature*, 415(6868), 141.
- Geider, R. J., & La Roche, J. (2002). Redfield revisited: variability of C [ratio] N [ratio] P in marine microalgae and its biochemical basis. *European Journal of Phycology*, 37(1), 1-17.
- Georgianna, D. R., & Mayfield, S. P. (2012). Exploiting diversity and synthetic biology for the production of algal biofuels. *Nature*, 488(7411), 329.
- Ghaffour, N., Bundschuh, J., Mahmoudi, H., & Goosen, M. F. (2015). Renewable energy-driven desalination technologies: A comprehensive review on challenges and potential applications of integrated systems. *Desalination*, 356, 94-114.
- Ghafoor, A., Hay, I. D., & Rehm, B. H. (2011). Role of exopolysaccharides in *Pseudomonas aeruginosa* biofilm formation and architecture. *Applied and environmental microbiology*, 77(15), 5238-5246.
- Gieger, C., Geistlinger, L., Altmaier, E., De Angelis, M. H., Kronenberg, F., Meitinger, T., Adamski, J. (2008). Genetics meets metabolomics: a genome-wide association study of metabolite profiles in human serum. *PLoS genetics*, 4(11), e1000282.
- Girard, J.-M., Roy, M.-L., Hafsa, M. B., Gagnon, J., Faucheux, N., Heitz, M., Deschênes, J.-S. (2014). Mixotrophic cultivation of green microalgae *Scenedesmus obliquus* on cheese whey permeate for biodiesel production. *Algal Research*, 5, 241-248.
- Glemser, M., Heining, M., Schmidt, J., Becker, A., Garbe, D., Buchholz, R., & Brück, T. (2016). Application of light-emitting diodes (LEDs) in cultivation of

- phototrophic microalgae: current state and perspectives. *Applied microbiology and biotechnology*, 100(3), 1077-1088.
- Goiris, K., Muylaert, K., Fraeye, I., Foubert, I., De Brabanter, J., & De Cooman, L. (2012). Antioxidant potential of microalgae in relation to their phenolic and carotenoid content. *Journal of Applied Phycology*, 24(6), 1477-1486.
- Goldemberg, J. (2007). Ethanol for a sustainable energy future. *Science*, 315(5813), 808-810.
- Gomez-Arreaza, A., Acosta, H., Quinones, W., Concepción, J. L., Michels, P. A., & Avilan, L. (2014). Extracellular functions of glycolytic enzymes of parasites: unpredicted use of ancient proteins. *Molecular and biochemical parasitology*, 193(2), 75-81.
- Gong, Y., Hu, H., Gao, Y., Xu, X., & Gao, H. (2011). Microalgae as platforms for production of recombinant proteins and valuable compounds: progress and prospects. *Journal of industrial microbiology & biotechnology*, 38(12), 1879-1890.
- González-Fernández, C., & Ballesteros, M. (2012). Linking microalgae and cyanobacteria culture conditions and key-enzymes for carbohydrate accumulation. *Biotechnology advances*, 30(6), 1655-1661.
- Gordillo, F. J., Jiménez, C., Figueroa, F. L., & Niell, F. X. (1998). Effects of increased atmospheric CO₂ and N supply on photosynthesis, growth and cell composition of the cyanobacterium *Spirulina platensis* (Arthrospira). *Journal of Applied Phycology*, 10(5), 461.
- Görg, A., Obermaier, C., Boguth, G., Harder, A., Scheibe, B., Wildgruber, R., & Weiss, W. (2000). The current state of two-dimensional electrophoresis with immobilized pH gradients. *ELECTROPHORESIS: An International Journal*, 21(6), 1037-1053.
- Görg, A., Postel, W., & Günther, S. (1988). Two-dimensional electrophoresis. The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*, 9(9), 531-546.
- Gouveia, J. D., Ruiz, J., van den Broek, L. A., Hesselink, T., Peters, S., Kleinegris, D. M., . . . Wijffels, R. H. (2017). *Botryococcus braunii* strains compared for biomass productivity, hydrocarbon and carbohydrate content. *Journal of biotechnology*, 248, 77-86.
- Gray, H. B. (2009). Powering the planet with solar fuel. *Nature chemistry*, 1(1), 7.

- Gremse, M., Chang, A., Schomburg, I., Grote, A., Scheer, M., Ebeling, C., & Schomburg, D. (2010). The BRENDA Tissue Ontology (BTO): the first all-integrating ontology of all organisms for enzyme sources. *Nucleic acids research*, 39(suppl_1), D507-D513.
- Grima, E. M., Belarbi, E.-H., Fernández, F. A., Medina, A. R., & Chisti, Y. (2003). Recovery of microalgal biomass and metabolites: process options and economics. *Biotechnology advances*, 20(7-8), 491-515.
- Grobbelaar, J. U. (2012). Microalgae mass culture: the constraints of scaling-up. *Journal of Applied Phycology*, 24(3), 315-318.
- Groombridge, B., Jenkins, M. D., & Jenkins, M. (2002). *World atlas of biodiversity: earth's living resources in the 21st century*: Univ of California Press.
- Guevara, M., Arredondo-Vega, B. O., Palacios, Y., Saéz, K., & Gómez, P. I. (2016). Comparison of growth and biochemical parameters of two strains of *Rhodomonas salina* (Cryptophyceae) cultivated under different combinations of irradiance, temperature, and nutrients. *Journal of Applied Phycology*, 28(5), 2651-2660.
- Gundry, R. L., White, M. Y., Murray, C. I., Kane, L. A., Fu, Q., Stanley, B. A., & Van Eyk, J. E. (2010). Preparation of proteins and peptides for mass spectrometry analysis in a bottom-up proteomics workflow. *Current protocols in molecular biology*, 90(1), 10.25. 11-10.25. 23.
- Gupta, V., Natarajan, C., Kumar, K., & Prasanna, R. (2011). Identification and characterization of endoglucanases for fungicidal activity in *Anabaena laxa* (Cyanobacteria). *Journal of Applied Phycology*, 23(1), 73-81.
- Guschina, I. A., & Harwood, J. L. (2006). Lipids and lipid metabolism in eukaryotic algae. *Progress in lipid research*, 45(2), 160-186.
- Hajjaj, H., Blanc, P., Goma, G., & Francois, J. (1998). Sampling techniques and comparative extraction procedures for quantitative determination of intra-and extracellular metabolites in filamentous fungi. *FEMS microbiology letters*, 164(1), 195-200.
- Halim, R., Danquah, M. K., & Webley, P. A. (2012). Extraction of oil from microalgae for biodiesel production: a review. *Biotechnology advances*, 30(3), 709-732.
- Hanotu, J., Bandulasena, H., & Zimmerman, W. B. (2012). Microflotation performance for algal separation. *Biotechnol Bioeng*, 109(7), 1663-1673.

- Harper, J. T., & Keeling, P. J. (2004). Lateral gene transfer and the complex distribution of insertions in eukaryotic enolase. *Gene*, 340(2), 227-235.
- Harun, R., & Danquah, M. K. (2011). Enzymatic hydrolysis of microalgal biomass for bioethanol production. *Chemical Engineering Journal*, 168(3), 1079-1084.
- Harun, R., Danquah, M. K., & Forde, G. M. (2010). Microalgal biomass as a fermentation feedstock for bioethanol production. *Journal of chemical technology and biotechnology*, 85(2), 199-203.
- He, X., Dai, J., & Wu, Q. (2016). Identification of sporopollenin as the outer layer of cell wall in microalga *Chlorella protothecoides*. *Frontiers in microbiology*, 7, 1047.
- Hehemann, J.-H., Redecke, L., Perbandt, M., Saborowski, R., & Betzel, C. (2007). Crystallization and preliminary X-ray diffraction studies of trypsin-like proteases from the gastric fluid of the marine crab *Cancer pagurus*. *Acta Crystallographica Section F: Structural Biology and Crystallization Communications*, 63(3), 242-245.
- Hehl, R., & Bülow, L. (2014). AthaMap web tools for the analysis of transcriptional and posttranscriptional regulation of gene expression in *Arabidopsis thaliana*. *Plant circadian networks* (pp. 139-156): Springer.
- Herrmann, V., & Jüttner, F. (1977). Excretion products of algae: Identification of biogenic amines by gas-liquid chromatography and mass spectrometry of their trifluoroacetamides. *Analytical biochemistry*, 78(2), 365-373.
- Hershey, J. C., Hautmann, M., Thompson, M. M., Rothblum, L. I., Haystead, T. A., & Owens, G. K. (1995). Angiotensin II-induced hypertrophy of rat vascular smooth muscle is associated with increased 18 S rRNA synthesis and phosphorylation of the rRNA transcription factor, upstream binding factor. *Journal of biological chemistry*, 270(42), 25096-25101.
- Himmel, M. E., Ding, S.-Y., Johnson, D. K., Adney, W. S., Nimlos, M. R., Brady, J. W., & Foust, T. D. (2007). Biomass recalcitrance: engineering plants and enzymes for biofuels production. *Science*, 315(5813), 804-807.
- Hinsby, A. M., Olsen, J. V., Bennett, K. L., & Mann, M. (2003). Signaling initiated by overexpression of the fibroblast growth factor receptor-1 investigated by mass spectrometry. *Molecular & Cellular Proteomics*, 2(1), 29-36.

- Ho, S.-H., Chen, C.-Y., Lee, D.-J., & Chang, J.-S. (2011). Perspectives on microalgal CO₂-emission mitigation systems a review. *Biotechnology advances*, 29(2), 189-198.
- Ho, Y., Gruhler, A., Heilbut, A., Bader, G. D., Moore, L., Adams, S.-L., Boutilier, K. (2002). Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature*, 415(6868), 180.
- Hopkinson, B. M., Meile, C., & Shen, C. (2013). Quantification of extracellular carbonic anhydrase activity in two marine diatoms and investigation of its role. *Plant physiology*, 162(2), 1142-1152.
- Hu, Q. (2004). *Environmental effects on cell composition* (Vol. 1): Blackwell Science Ltd.: Oxford, UK.
- Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., & Darzins, A. (2008). Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *The Plant Journal*, 54(4), 621-639.
- Huber, G. W., Chheda, J. N., Barrett, C. J., & Dumesic, J. A. (2005). Production of liquid alkanes by aqueous-phase processing of biomass-derived carbohydrates. *Science*, 308(5727), 1446-1450.
- Huerlimann, R., De Nys, R., & Heimann, K. (2010). Growth, lipid content, productivity, and fatty acid composition of tropical microalgae for scale-up production. *Biotechnol Bioeng*, 107(2), 245-257.
- Huo, Y.-X., Cho, K. M., Rivera, J. G. L., Monte, E., Shen, C. R., Yan, Y., & Liao, J. C. (2011). Conversion of proteins into biofuels by engineering nitrogen flux. *Nature biotechnology*, 29(4), 346.
- Husain, Q. (2010). β Galactosidases and their potential applications: a review. *Critical reviews in biotechnology*, 30(1), 41-62.
- Idle, J. R., & Gonzalez, F. J. (2007). Metabolomics. *Cell metabolism*, 6(5), 348-351.
- Ike, M., Ko, Y., Yokoyama, K., Sumitani, J.-I., Kawaguchi, T., Ogasawara, W., Morikawa, Y. (2007). Cellobiohydrolase I (Cel7A) from *Trichoderma reesei* has chitosanase activity. *Journal of Molecular Catalysis B: Enzymatic*, 47(3-4), 159-163.
- Ioelovich, M. (2013). Energetic potential of plant biomass and its use. *Int. J. Renew. Sustain. Energy*, 2(2), 26.
- Ioelovich, M. (2015). Recent findings and the energetic potential of plant biomass as a renewable source of biofuels—a review. *Bioresources*, 10(1), 1879-1914.

- Ito, T., Tanaka, M., Shinkawa, H., Nakada, T., Ano, Y., Kurano, N., Tomita, M. (2013). Metabolic and morphological changes of an oil accumulating trebouxiphycean alga in nitrogen-deficient conditions. *Metabolomics*, 9(1), 178-187.
- James, G. O., Hocart, C. H., Hillier, W., Price, G. D., & Djordjevic, M. A. (2013). Temperature modulation of fatty acid profiles for biofuel production in nitrogen deprived *Chlamydomonas reinhardtii*. *Bioresource technology*, 127, 441-447.
- Jeon, S., Jeong, B., & Chang, Y. (2017). Consequences of microbial interactions with hydrocarbons, oils, and lipids: production of fuels and chemicals.
- Jeong, B.-r., Wu-Scharf, D., Zhang, C., & Cerutti, H. (2002). Suppressors of transcriptional transgenic silencing in *Chlamydomonas* are sensitive to DNA-damaging agents and reactivate transposable elements. *Proceedings of the National Academy of Sciences*, 99(2), 1076-1081.
- Jewett, M. C., Hofmann, G., & Nielsen, J. (2006). Fungal metabolite analysis in genomics and phenomics. *Current opinion in biotechnology*, 17(2), 191-197.
- John, R. P., Anisha, G., Nampoothiri, K. M., & Pandey, A. (2011). Micro and macroalgal biomass: a renewable source for bioethanol. *Bioresource technology*, 102(1), 186-193.
- Johnson, X., & Alric, J. (2013). Central carbon metabolism and electron transport in *Chlamydomonas reinhardtii*: metabolic constraints for carbon partitioning between oil and starch. *Eukaryotic cell*, 12(6), 776-793.
- Juneja, A., Ceballos, R. M., & Murthy, G. S. (2013). Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: a review. *Energies*, 6(9), 4607-4638.
- Kang, J., & Wen, Z. (2015). A manuscript accepted by Algal Research. *Environmental stewardship by microalgae: air and water*, 41.
- Kang, M., Abdelmageed, H., Lee, S., Reichert, A., Mysore, K. S., & Allen, R. D. (2013). At MBP-1, an alternative translation product of LOS 2, affects abscisic acid responses and is modulated by the E 3 ubiquitin ligase A t SAP 5. *The Plant Journal*, 76(3), 481-493.
- Katiyar, R., Gurjar, B., Biswas, S., Pruthi, V., Kumar, N., & Kumar, P. (2017). Microalgae: an emerging source of energy based bio-products and a solution

- for environmental issues. *Renewable and Sustainable Energy Reviews*, 72, 1083-1093.
- Katrolia, P., Rajashekhara, E., Yan, Q., & Jiang, Z. (2014). Biotechnological potential of microbial α -galactosidases. *Critical reviews in biotechnology*, 34(4), 307-317.
- Ketheesan, B., & Nirmalakhandan, N. (2012). Feasibility of microalgal cultivation in a pilot-scale airlift-driven raceway reactor. *Bioresource technology*, 108, 196-202.
- Khan, M. I., Shin, J. H., & Kim, J. D. (2018). The promising future of microalgae: current status, challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and other products. *Microbial cell factories*, 17(1), 36.
- Khoeiyi, Z. A., Seyfabadi, J., & Ramezanpour, Z. (2012). Effect of light intensity and photoperiod on biomass and fatty acid composition of the microalgae, *Chlorella vulgaris*. *Aquaculture International*, 20(1), 41-49.
- Kimura, S., & Kondo, T. (2002). Recent progress in cellulose biosynthesis. *Journal of plant research*, 115(4), 297-302.
- Kleinová, A., Cvengrošová, Z., Rimarčík, J., Buzetzki, E., Mikulec, J., & Cvengroš, J. (2012). Biofuels from algae. *Procedia Engineering*, 42, 231-238.
- Kligerman, D. C., & Bouwer, E. J. (2015). Prospects for biodiesel production from algae-based wastewater treatment in Brazil: A review. *Renewable and Sustainable Energy Reviews*, 52, 1834-1846.
- Kliphuis, A. M., Klok, A. J., Martens, D. E., Lamers, P. P., Janssen, M., & Wijffels, R. H. (2012). Metabolic modeling of *Chlamydomonas reinhardtii*: energy requirements for photoautotrophic growth and maintenance. *Journal of Applied Phycology*, 24(2), 253-266.
- Klose, J. (1975). Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues. *Humangenetik*, 26(3), 231-243.
- Knothe, G. (2005). Dependence of biodiesel fuel properties on the structure of fatty acid alkyl esters. *Fuel processing technology*, 86(10), 1059-1070.
- Knothe, G. (2008). "Designer" biodiesel: optimizing fatty ester composition to improve fuel properties. *Energy & Fuels*, 22(2), 1358-1364.

- Komárek, J., & Marvan, P. (1992). Morphological differences in natural populations of the genus *Botryococcus* (Chlorophyceae). *Archiv für Protistenkunde*, 141(1-2), 65-100.
- Kong, Q.-x., Li, L., Martinez, B., Chen, P., & Ruan, R. (2010). Culture of microalgae *Chlamydomonas reinhardtii* in wastewater for biomass feedstock production. *Applied biochemistry and biotechnology*, 160(1), 9.
- Kotwal, S., Gote, M., Sainkar, S., Khan, M., & Khire, J. (1998). Production of α -galactosidase by thermophilic fungus *Humicola* sp. in solid-state fermentation and its application in soyamilk hydrolysis. *Process Biochemistry*, 33(3), 337-343.
- Kromkamp, J. (1987). Formation and functional significance of storage products in cyanobacteria. *New Zealand Journal of Marine and Freshwater Research*, 21(3), 457-465.
- Krzemińska, I., Pawlik-Skowrońska, B., Trzcińska, M., & Tys, J. (2014). Influence of photoperiods on the growth rate and biomass productivity of green microalgae. *Bioprocess and biosystems engineering*, 37(4), 735-741.
- Kumar, A., Ergas, S., Yuan, X., Sahu, A., Zhang, Q., Dewulf, J., . . . Van Langenhove, H. (2010). Enhanced CO₂ fixation and biofuel production via microalgae: recent developments and future directions. *Trends in biotechnology*, 28(7), 371-380.
- Kumar, M., Sharma, M., & Dwivedi, G. (2013). Algae oil as future energy source in Indian perspective. *International Journal of Renewable Energy Research (IJRER)*, 3(4), 913-921.
- Kurien, B. T., & Scofield, R. H. (2012). *Protein electrophoresis: methods and protocols*: Springer.
- Kurinjimalar, C., Kavitha, G., Shamshath-Begum, S., Rajaram, G., Nagaraj, S., Senthilkumar, N., & Rengasamy, R. (2017). Optimization and production of *Botryococcus braunii* biomass using commercial nutrients by Response Surface Methodology. *Iranica Journal of Energy and Environment*, 8(1), 18-25.
- La Roche, J., Geider, R. J., Graziano, L. M., Murray, H., & Lewis, K. (1993). Induction of specific proteins in eukaryotic algae grown under iron, phosphorus, or nitrogen deficient conditions 1. *Journal of phycology*, 29(6), 767-777.

- Laemmli, J. (1970a). SDS-polyacrylamide gel electrophoresis of protein. *Nature*, 2, 680-685.
- Laemmli, U. K. (1970b). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227(5259), 680.
- Lane, A. E., & Burris, J. E. (1981). Effects of environmental pH on the internal pH of *Chlorella pyrenoidosa*, *Scenedesmus quadricauda*, and *Euglena mutabilis*. *Plant physiology*, 68(2), 439-442.
- Largeau, C., Casadevall, E., Berkloff, C., & Dhamelincourt, P. (1980). Sites of accumulation and composition of hydrocarbons in *Botryococcus braunii*. *Phytochemistry*, 19(6), 1043-1051.
- Latiffi, N. A. A., Mohamed, R. M. S. R., Apandi, N. M., & Tajuddin, R. M. (2017). *Preliminary Assessment of Growth Rates on Different Concentration of Microalgae Scenedesmus sp. in Industrial Meat Food Processing Wastewater*. Paper presented at the MATEC Web of Conferences.
- Lee, C.-H., Chae, H.-S., Lee, S.-H., & Kim, H. S. (2015). Growth characteristics and lipid content of three Korean isolates of *Botryococcus braunii* (Trebouxiophyceae). *Journal of Ecology and Environment*, 38(1), 67-74.
- Lee, O. K., Kim, A. L., Seong, D. H., Lee, C. G., Jung, Y. T., Lee, J. W., & Lee, E. Y. (2013). Chemo-enzymatic saccharification and bioethanol fermentation of lipid-extracted residual biomass of the microalga, *Dunaliella tertiolecta*. *Bioresource technology*, 132, 197-201.
- Leonardos, N., & Geider, R. J. (2004). Responses of elemental and biochemical composition of *Chaetoceros muelleri* to growth under varying light and nitrate: phosphate supply ratios and their influence on critical N: P. *Limnology and Oceanography*, 49(6), 2105-2114.
- Lewis, T. S., Hunt, J. B., Aveline, L. D., Jonscher, K. R., Louie, D. F., Yeh, J. M., Ahn, N. G. (2000). Identification of novel MAP kinase pathway signaling targets by functional proteomics and mass spectrometry. *Molecular cell*, 6(6), 1343-1354.
- Li, T., Zheng, Y., Yu, L., & Chen, S. (2013a). High productivity cultivation of a heat-resistant microalga *Chlorella sorokiniana* for biofuel production. *Bioresource technology*, 131, 60-67.
- Li, W. (1997). *Molecular evolution*: Sinauer associates incorporated.

- Li, X., Příbyl, P., Bišová, K., Kawano, S., Cepák, V., Zachleder, V., Vítová, M. (2013b). The microalga *Parachlorella kessleri*—A novel highly efficient lipid producer. *Biotechnol Bioeng*, 110(1), 97-107.
- Li, Y., Han, D., Hu, G., Sommerfeld, M., & Hu, Q. (2010). Inhibition of starch synthesis results in overproduction of lipids in *Chlamydomonas reinhardtii*. *Biotechnol Bioeng*, 107(2), 258-268.
- Li, Y., Han, D., Sommerfeld, M., & Hu, Q. (2011). Photosynthetic carbon partitioning and lipid production in the oleaginous microalga *Pseudochlorococcum* sp.(Chlorophyceae) under nitrogen-limited conditions. *Bioresource technology*, 102(1), 123-129.
- Li, Y., Horsman, M., Wang, B., Wu, N., & Lan, C. Q. (2008a). Effects of nitrogen sources on cell growth and lipid accumulation of green alga *Neochloris oleoabundans*. *Applied microbiology and biotechnology*, 81(4), 629-636.
- Li, Y., Horsman, M., Wu, N., Lan, C. Q., & Dubois-Calero, N. (2008b). Biofuels from microalgae. *Biotechnology progress*, 24(4), 815-820.
- Lilley, K. S., & Friedman, D. B. (2004). All about DIGE: quantification technology for differential-display 2D-gel proteomics. *Expert review of proteomics*, 1(4), 401-409.
- Lin, C.-Y., & Li, R.-J. (2009). Fuel properties of biodiesel produced from the crude fish oil from the soapstock of marine fish. *Fuel processing technology*, 90(1), 130-136.
- Liu, J., Huang, J., Sun, Z., Zhong, Y., Jiang, Y., & Chen, F. (2011). Differential lipid and fatty acid profiles of photoautotrophic and heterotrophic *Chlorella zofingiensis*: assessment of algal oils for biodiesel production. *Bioresource technology*, 102(1), 106-110.
- Liu, J., & Xia, W. (2005). Characterization of chitosanase from cellulase produced by *Trichoderma viride*. *CHINESE JOURNAL OF BIOCHEMISTRY AND MOLECULAR BIOLOGY*, 21(5), 713.
- Liu, L., Pohnert, G., & Wei, D. (2016). Extracellular metabolites from industrial microalgae and their biotechnological potential. *Marine drugs*, 14(10), 191.
- Liu, S., Zhao, Y., Liu, L., Ao, X., Ma, L., Wu, M., & Ma, F. (2015). Improving cell growth and lipid accumulation in green microalgae *Chlorella* sp. Via UV irradiation. *Applied biochemistry and biotechnology*, 175(7), 3507-3518.

- Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *Journal of biological chemistry*, 193(1), 265-275.
- Mahendran, S., Saravanan, S., Vijayabaskar, P., Anandapandian, K., & Shankar, T. (2013). Antibacterial potential of microbial exopolysaccharide from *Ganoderma lucidum* and *Lysinibacillus fusiformis*. *Int J Recent Sci Res*, 4, 501-505.
- Maity, J. P., Bundschuh, J., Chen, C.-Y., & Bhattacharya, P. (2014). Microalgae for third generation biofuel production, mitigation of greenhouse gas emissions and wastewater treatment: Present and future perspectives—A mini review. *Energy*, 78, 104-113.
- Marcilla, A., Gómez-Siurana, A., & Menargues, S. (2005a). Qualitative study of the evolution of the composition of the gas evolved in the thermal and HY-catalytic oxidative degradation of EVA copolymers. *Thermochimica acta*, 438(1-2), 155-163.
- Marcilla, A., Gómez, A., & Menargues, S. (2005b). TGA/FTIR study of the catalytic pyrolysis of ethylene–vinyl acetate copolymers in the presence of MCM-41. *Polymer degradation and stability*, 89(1), 145-152.
- Markou, G., Angelidaki, I., & Georgakakis, D. (2012). Microalgal carbohydrates: an overview of the factors influencing carbohydrates production, and of main bioconversion technologies for production of biofuels. *Applied microbiology and biotechnology*, 96(3), 631-645.
- Markou, G., & Georgakakis, D. (2011). Cultivation of filamentous cyanobacteria (blue-green algae) in agro-industrial wastes and wastewaters: a review. *Applied energy*, 88(10), 3389-3401.
- Martin, M., Biver, S., Steels, S., Barbeyron, T., Jam, M., Portetelle, D., . . . Vandenbol, M. (2014). Functional screening of a metagenomic library of seaweed-associated microbiota: identification and characterization of a halotolerant, cold-active marine endo- β -1, 4-endoglucanase. *Applied and environmental microbiology*, AEM. 01194-01114.
- Mata, T. M., Martins, A. A., & Caetano, N. S. (2010). Microalgae for biodiesel production and other applications: a review. *Renewable and Sustainable Energy Reviews*, 14(1), 217-232.

- Mathur, R., Reddy, M., Yee, S., Imbesi, R., Groth-Vasselli, B., & Farnsworth, P. (1992). Investigation of lens glycolytic enzymes: species distribution and interaction with supramolecular order. *Experimental eye research*, 54(2), 253-260.
- McCourt, R. M., Delwiche, C. F., & Karol, K. G. (2004). Charophyte algae and land plant origins. *Trends in Ecology & Evolution*, 19(12), 661-666.
- McKay, R. M. L., La Roche, J., Yakunin, A. F., Durnford, D. G., & Geider, R. J. (1999). Accumulation of ferredoxin and flavodoxin in a marine diatom in response to Fe. *Journal of phycology*, 35(3), 510-519.
- Melis, A. (2013). Carbon partitioning in photosynthesis. *Current Opinion in Chemical Biology*, 17(3), 453-456.
- Menichelli, C., Gascuel, O., & Bréhélin, L. (2018). Improving pairwise comparison of protein sequences with domain co-occurrence. *PLoS computational biology*, 14(1), e1005889.
- Metting, F. (1996). Biodiversity and application of microalgae. *Journal of industrial microbiology*, 17(5-6), 477-489.
- Metz, B., Davidson, O. R., Bosch, P. R., Dave, R., & Meyer, L. A. (2007). Contribution of working group III to the fourth assessment report of the intergovernmental panel on climate change, 2007: Cambridge University Press, Cambridge, UK.
- Metzger, P., Berkaloﬀ, C., Casadevall, E., & Coute, A. (1985). Alkadiene-and botryococcene-producing races of wild strains of *Botryococcus braunii*. *Phytochemistry*, 24(10), 2305-2312.
- Miao, X., Wu, Q., & Yang, C. (2004). Fast pyrolysis of microalgae to produce renewable fuels. *Journal of analytical and applied pyrolysis*, 71(2), 855-863.
- Michel, G., & Czjzek, M. (2013). Polysaccharide-degrading enzymes from marine bacteria *Marine Enzymes for Biocatalysis* (pp. 429-464): Elsevier.
- Milano, J., Ong, H. C., Masjuki, H., Chong, W., Lam, M. K., Loh, P. K., & Vellayan, V. (2016). Microalgae biofuels as an alternative to fossil fuel for power generation. *Renewable and Sustainable Energy Reviews*, 58, 180-197.
- Miller, R., Wu, G., Deshpande, R. R., Vieler, A., Gärtner, K., Li, X., Liu, B. (2010). Changes in transcript abundance in *Chlamydomonas reinhardtii* following nitrogen-deprivation predict diversion of metabolism. *Plant physiology*, pp. 110.165159.

- Mishra, N. C. (2010). *Introduction to proteomics: principles and applications* (Vol. 146): Wiley Online Library.
- Moellering, E. R., & Benning, C. (2010). RNA interference silencing of a major lipid droplet protein affects lipid droplet size in *Chlamydomonas reinhardtii*. *Eukaryotic cell*, 9(1), 97-106.
- Mohammed, K. (2013). Microalgal photobioreactors for carbon-efficient wastewater treatment.
- Mohan, S. V., Rohit, M., Chiranjeevi, P., Chandra, R., & Navaneeth, B. (2015). Heterotrophic microalgae cultivation to synergize biodiesel production with waste remediation: progress and perspectives. *Bioresource technology*, 184, 169-178.
- Molnár, I., Lopez, D., Wisecaver, J. H., Devarenne, T. P., Weiss, T. L., Pellegrini, M., & Hackett, J. D. (2012). Bio-crude transcriptomics: gene discovery and metabolic network reconstruction for the biosynthesis of the terpenome of the hydrocarbon oil-producing green alga, *Botryococcus braunii* race B (Showa). *BMC genomics*, 13(1), 576.
- Morales, M., Sánchez, L., & Revah, S. (2017). The impact of environmental factors on CO₂ fixation by microalgae. *FEMS microbiology letters*.
- Mori, N., Moriyama, T., & Sato, N. (2019). Uncommon properties of lipid biosynthesis of isolated plastids in the unicellular red alga *Cyanidioschyzon merolae*. *FEBS open bio*, 9(1), 114-128.
- Moroney, J. V. (2001). Algal photosynthesis. *e LS*.
- Moss, B. (1973). The influence of environmental factors on the distribution of freshwater algae: an experimental study: II. The role of pH and the carbon dioxide-bicarbonate system. *The Journal of Ecology*, 157-177.
- Mou, S., Zhang, Y., Li, G., Li, H., Liang, Y., Tang, L., Zhang, C. (2017). Effects of elevated CO₂ and nitrogen supply on the growth and photosynthetic physiology of a marine cyanobacterium, *Synechococcus* sp. PCC7002. *Journal of Applied Phycology*, 29(4), 1755-1763.
- Msanne, J., Xu, D., Konda, A. R., Casas-Mollano, J. A., Awada, T., Cahoon, E. B., & Cerutti, H. (2012). Metabolic and gene expression changes triggered by nitrogen deprivation in the photoautotrophically grown microalgae *Chlamydomonas reinhardtii* and *Coccomyxa* sp. C-169. *Phytochemistry*, 75, 50-59.

- Mudimu, O., Rybalka, N., Bauersachs, T., Friedl, T., & Schulz, R. (2015). Influence of different CO₂ concentrations on microalgae growth, α -tocopherol content and fatty acid composition. *Geomicrobiology Journal*, 32(3-4), 291-303.
- Munir, N., Imtiaz, A., Sharif, N., & Naz, S. (2015). Optimization of growth conditions of different algal strains and determination of their lipid contents. *J Anim Plant Sci*, 25(2), 546-553.
- Muradyan, E., Klyachko-Gurvich, G., Tsoglin, L., Sergeyenko, T., & Pronina, N. (2004). Changes in lipid metabolism during adaptation of the *Dunaliella salina* photosynthetic apparatus to high CO₂ concentration. *Russian journal of plant physiology*, 51(1), 53-62.
- Nahak, S., Nahak, G., Pradhan, I., & Sahu, R. (2011). Bioethanol from marine algae: a solution to global warming problem. *J. Appl. Environ. Biol. Sci*, 1(4), 74-80.
- Naito, K., Matsui, M., & Imai, I. (2005). Ability of marine eukaryotic red tide microalgae to utilize insoluble iron. *Harmful Algae*, 4(6), 1021-1032.
- Nascimento, I. A., Marques, S. S. I., Cabanelas, I. T. D., Pereira, S. A., Druzian, J. I., de Souza, C. O., Nascimento, M. A. (2013). Screening microalgae strains for biodiesel production: lipid productivity and estimation of fuel quality based on fatty acids profiles as selective criteria. *Bioenergy research*, 6(1), 1-13.
- Natrah, F., Yusoff, F., Shariff, M., Abas, F., & Mariana, N. (2007). Screening of Malaysian indigenous microalgae for antioxidant properties and nutritional value. *Journal of Applied Phycology*, 19(6), 711-718.
- Nguyen, D., & Nguyen, T. (2014). Detection on antioxidant and cytotoxicity activities of exopolysaccharides isolated in plant-originated *Lactococcus lactis*. *Biomedical & Pharmacology Journal*, 7(1), 33-38.
- Nichols, H. W. (1973). Growth media-freshwater. *Culture methods and growth measurements*, 7-24.
- Nichols, H. W., & Bold, H. C. (1965). *Trichosarcina polymorpha* gen. et sp. nov. *Journal of phycology*, 1(1), 34-38.
- Noctor, G., Veljovic-Jovanovic, S., Driscoll, S., Novitskaya, L., & Foyer, C. H. (2002). Drought and oxidative load in the leaves of C3 plants: a predominant role for photorespiration? *Annals of Botany*, 89(7), 841-850.
- O'Farrell, P. H. (1975). High resolution two-dimensional electrophoresis of proteins. *Journal of biological chemistry*, 250(10), 4007-4021.

- Ohlrogge, J. B., & Jaworski, J. G. (1997). Regulation of fatty acid synthesis. *Annual review of plant biology*, 48(1), 109-136.
- Okada, S., Devarenne, T. P., & Chappell, J. (2000). Molecular characterization of squalene synthase from the green microalga *Botryococcus braunii*, race B. *Archives of biochemistry and biophysics*, 373(2), 307-317.
- Oliveira, C., Guimarães, P. M., & Domingues, L. (2011). Recombinant microbial systems for improved β -galactosidase production and biotechnological applications. *Biotechnology advances*, 29(6), 600-609.
- Oswald, W. J., & Gotaas, H. B. (1957). Photosynthesis in sewage treatment. *Trans. Am. Soc. Civ. Eng.*, 122(1), 73-105.
- Pandey, A., Fernandez, M. M., Steen, H., Blagoev, B., Nielsen, M. M., Roche, S., . . . Lodish, H. F. (2000). Identification of a novel immunoreceptor tyrosine-based activation motif-containing molecule, STAM2, by mass spectrometry and its involvement in growth factor and cytokine receptor signaling pathways. *Journal of biological chemistry*, 275(49), 38633-38639.
- Pane, L., Franceschi, E., De Nuccio, L., & Carli, A. (2001). Applications of thermal analysis on the marine phytoplankton, *Tetraselmis suecica*. *Journal of thermal analysis and calorimetry*, 66(1), 145-154.
- Patel, A., & Prajapat, J. (2013). Food and health applications of exopolysaccharides produced by lactic acid bacteria. *Advances in Dairy Research*, 1-8.
- Peng, W., Wu, Q., Tu, P., & Zhao, N. (2001). Pyrolytic characteristics of microalgae as renewable energy source determined by thermogravimetric analysis. *Bioresource technology*, 80(1), 1-7.
- Petroleum, B. (2012). BP statistical review of world energy: London: British Petroleum.
- Pillai, S. A., Chobisa, D., Urimi, D., & Ravindra, N. (2016). Pharmaceutical Glass Interactions: A Review of Possibilities. *J Pharm Sci Res*, 8, 103-111.
- Placzek, S., Schomburg, I., Chang, A., Jeske, L., Ulbrich, M., Tillack, J., & Schomburg, D. (2016). BRENDA in 2017: new perspectives and new tools in BRENDA. *Nucleic acids research*, gkw952.
- Plieth, C., Sattelmacher, B., & Hansen, U.-P. (1997). Cytoplasmic Ca^{2+} - H^{+} -exchange buffers in green algae. *Protoplasma*, 198(1-2), 107-124.

- Polle, J. E., Neofotis, P., Huang, A., Chang, W., Sury, K., & Wiech, E. M. (2014). Carbon partitioning in green algae (chlorophyta) and the enolase enzyme. *Metabolites*, 4(3), 612-628.
- Powell, E., & Hill, G. (2009). Economic assessment of an integrated bioethanol–biodiesel–microbial fuel cell facility utilizing yeast and photosynthetic algae. *Chemical Engineering Research and Design*, 87(9), 1340-1348.
- Prabhakar, V., Löttgert, T., Geimer, S., Dörmann, P., Krüger, S., Vijayakumar, V., . . . Feussner, I. (2010). Phosphoenolpyruvate provision to plastids is essential for gametophyte and sporophyte development in *Arabidopsis thaliana*. *The Plant Cell*, 22(8), 2594-2617.
- Prabhakar, V., Löttgert, T., Gigolashvili, T., Bell, K., Flügge, U.-I., & Häusler, R. E. (2009). Molecular and functional characterization of the plastid-localized phosphoenolpyruvate enolase (ENO1) from *Arabidopsis thaliana*. *FEBS letters*, 583(6), 983-991.
- Prescott, C. (1978a). CY Prescott et al., Phys. Lett. 77B, 347 (1978). *Phys. Lett.*, 77, 347.
- Prescott, G. (1978b). How to know freshwater algae. 3rd Edn., Wes. C: Brown Company Publishers Dubuque, IOWA.
- Pulz, O., & Gross, W. (2004). Valuable products from biotechnology of microalgae. *Applied microbiology and biotechnology*, 65(6), 635-648.
- Qin, J. G., & Li, Y. (2006). Optimization of the growth environment of *Botryococcus braunii* strain CHN 357. *Journal of Freshwater Ecology*, 21(1), 169-176.
- Radakovits, R., Eduafo, P. M., & Posewitz, M. C. (2011). Genetic engineering of fatty acid chain length in *Phaeodactylum tricornutum*. *Metabolic engineering*, 13(1), 89-95.
- Radakovits, R., Jinkerson, R. E., Darzins, A., & Posewitz, M. C. (2010). Genetic engineering of algae for enhanced biofuel production. *Eukaryotic cell*, 9(4), 486-501.
- Ramazanov, A., & Ramazanov, Z. (2006). Isolation and characterization of a starchless mutant of *Chlorella pyrenoidosa* STL-PI with a high growth rate, and high protein and polyunsaturated fatty acid content. *Phycological Research*, 54(4), 255-259.

- Rao, A. R., Sarada, R., Baskaran, V., & Ravishankar, G. A. (2006). Antioxidant activity of *Botryococcus braunii* extract elucidated in vitro models. *Journal of agricultural and food chemistry*, 54(13), 4593-4599.
- Ratledge, C., & Cohen, Z. (2008). Microbial and algal oils: do they have a future for biodiesel or as commodity oils? *Lipid Technology*, 20(7), 155-160.
- Rawat, I., Kumar, R. R., Mutanda, T., & Bux, F. (2013). Biodiesel from microalgae: a critical evaluation from laboratory to large scale production. *Applied energy*, 103, 444-467.
- Reboloso-Fuentes, M. M., Navarro-Pérez, A., García-Camacho, F., Ramos-Miras, J., & Guil-Guerrero, J. (2001). Biomass nutrient profiles of the microalga *Nannochloropsis*. *Journal of agricultural and food chemistry*, 49(6), 2966-2972.
- Recht, L., Zarka, A., & Boussiba, S. (2012). Patterns of carbohydrate and fatty acid changes under nitrogen starvation in the microalgae *Haematococcus pluvialis* and *Nannochloropsis* sp. *Applied microbiology and biotechnology*, 94(6), 1495-1503.
- Renaud, S. M., Thinh, L.-V., Lambrinidis, G., & Parry, D. L. (2002). Effect of temperature on growth, chemical composition and fatty acid composition of tropical Australian microalgae grown in batch cultures. *Aquaculture*, 211(1-4), 195-214.
- Richardson, J. W., Johnson, M. D., & Outlaw, J. L. (2012). Economic comparison of open pond raceways to photo bio-reactors for profitable production of algae for transportation fuels in the Southwest. *Algal Research*, 1(1), 93-100.
- Richmond, A. (2008). *Handbook of microalgal culture: biotechnology and applied phycology*: John Wiley & Sons.
- Rier, S. T., Nawrocki, K. S., & Whitley, J. C. (2011). Response of biofilm extracellular enzymes along a stream nutrient enrichment gradient in an agricultural region of north central Pennsylvania, USA. *Hydrobiologia*, 669(1), 119-131.
- Rismani-Yazdi, H., Haznedaroglu, B. Z., Bibby, K., & Peccia, J. (2011). Transcriptome sequencing and annotation of the microalgae *Dunaliella tertiolecta*: pathway description and gene discovery for production of next-generation biofuels. *BMC genomics*, 12(1), 148.
- Rodolfi, L., Chini Zittelli, G., Bassi, N., Padovani, G., Biondi, N., Bonini, G., & Tredici, M. R. (2009). Microalgae for oil: Strain selection, induction of lipid

- synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnol Bioeng*, 102(1), 100-112.
- Roessner, U., & Beckles, D. M. (2009). Metabolite measurements *Plant metabolic networks* (pp. 39-69): Springer.
- Romani, A., & Sabater, S. (2000). Influence of algal biomass on extracellular enzyme activity in river biofilms. *Microbial Ecology*, 40(1), 16-24.
- Rosenberg, Z. M.-M. (2006). Current trends of β -galactosidase application in food technology. *J Food Nutr Res*, 45, 47-54.
- Ruan, K., Duan, J., Bai, F., Lemaire, M., Ma, X., & Bai, L. (2009). Function of *Dunaliella salina* (Dunaliellaceae) enolase and its expression during stress. *European Journal of Phycology*, 44(2), 207-214.
- Ruangsomboon, S. (2015). Effects of different media and nitrogen sources and levels on growth and lipid of green microalga *Botryococcus braunii* KMITL and its biodiesel properties based on fatty acid composition. *Bioresource technology*, 191, 377-384.
- Ruas-Madiedo, P., Hugenholtz, J., & Zoon, P. (2002). An overview of the functionality of exopolysaccharides produced by lactic acid bacteria. *International Dairy Journal*, 12(2-3), 163-171.
- Rubio, F. C., Camacho, F. G., Sevilla, J., Chisti, Y., & Grima, E. M. (2003). A mechanistic model of photosynthesis in microalgae. *Biotechnol Bioeng*, 81(4), 459-473.
- Rutschow, H., Ytterberg, A. J., Friso, G., Nilsson, R., & van Wijk, K. J. (2008). Quantitative proteomics of a chloroplast SRP54 sorting mutant and its genetic interactions with CLPC1 in Arabidopsis. *Plant physiology*, 148(1), 156-175.
- Saifullah, A., Karim, M. A., & Ahmad-Yazid, A. (2014). Microalgae: An alternative source of renewable energy. *Am. J. Eng. Res*, 3(3), 330-338.
- Sanada, C. T., Karp, S. G., Spier, M. R., Portella, A. C., Gouvêa, P. M., Yamaguishi, C. T., Soccol, C. R. (2009). Utilization of soybean vinasse for α -galactosidase production. *Food research international*, 42(4), 476-483.
- Sankaran, B., & Thiruneelagandan, E. (2015). Microalgal diversity of Parthasarathy temple tank, Chennai, India. *Int. J. Curr. Microbiol. App. Sci*, 4(4), 168-173.
- Sato, N., Hagio, M., Wada, H., & Tsuzuki, M. (2000). Environmental effects on acidic lipids of thylakoid membranes: Portland Press Limited.

- Sawayama, S., Minowa, T., & Yokoyama, S.-Y. (1999). Possibility of renewable energy production and CO₂ mitigation by thermochemical liquefaction of microalgae. *Biomass and Bioenergy*, 17(1), 33-39.
- Schenk, P. M., Thomas-Hall, S. R., Stephens, E., Marx, U. C., Mussgnug, J. H., Posten, C., Hankamer, B. (2008). Second generation biofuels: high-efficiency microalgae for biodiesel production. *Bioenergy research*, 1(1), 20-43.
- Schomburg, I., Jeske, L., Ulbrich, M., Placzek, S., Chang, A., & Schomburg, D. (2017). The BRENDA enzyme information system—From a database to an expert system. *Journal of biotechnology*, 261, 194-206.
- Schwenzfeier, A., Wierenga, P. A., & Gruppen, H. (2011). Isolation and characterization of soluble protein from the green microalgae *Tetraselmis* sp. *Bioresource technology*, 102(19), 9121-9127.
- Senousy, H. H., Beakes, G. W., & Hack, E. (2004). Phylogenetic placement of *Botryococcus braunii* (trebouxiophyceae) and *Botryococcus sudeticus* isolate utex 2629 (Chlorophyceae) 1. *Journal of phycology*, 40(2), 412-423.
- Sforza, E., Gris, B., de Farias Silva, C., Morosinotto, T., & Bertucco, A. (2014). Effects of light on cultivation of *Scenedesmus obliquus* in batch and continuous flat plate photobioreactor. *Chem. Eng*, 38, 211-216.
- Shen, C., & Hopkinson, B. M. (2015). Size scaling of extracellular carbonic anhydrase activity in centric marine diatoms. *Journal of phycology*, 51(2), 255-263.
- Shen, Y. (2014). Carbon dioxide bio-fixation and wastewater treatment via algae photochemical synthesis for biofuels production. *RSC Advances*, 4(91), 49672-49722.
- Shen, Y., Zhang, H., Xu, X., & Lin, X. (2015). Biofilm formation and lipid accumulation of attached culture of *Botryococcus braunii*. *Bioprocess and biosystems engineering*, 38(3), 481-488.
- Sheng, J., Kim, H. W., Badalamenti, J. P., Zhou, C., Sridharakrishnan, S., Krajmalnik-Brown, R., Vannela, R. (2011). Effects of temperature shifts on growth rate and lipid characteristics of *Synechocystis* sp. PCC6803 in a bench-top photobioreactor. *Bioresource technology*, 102(24), 11218-11225.
- Sibi, G., Shetty, V., & Mokashi, K. (2016). Enhanced lipid productivity approaches in microalgae as an alternate for fossil fuels—A review. *Journal of the Energy Institute*, 89(3), 330-334.

- Singh, S., & Singh, P. (2015). Effect of temperature and light on the growth of algae species: a review. *Renewable and Sustainable Energy Reviews*, 50, 431-444.
- Smith, B. J. (1994). SDS polyacrylamide gel electrophoresis of proteins *Basic protein and peptide protocols* (pp. 23-34): Springer.
- Smith, J. B. (2001). Peptide sequencing by Edman degradation. *e LS*.
- Solovchenko, A., Khozin-Goldberg, I., Didi-Cohen, S., Cohen, Z., & Merzlyak, M. (2008). Effects of light intensity and nitrogen starvation on growth, total fatty acids and arachidonic acid in the green microalga *Parietochloris incisa*. *Journal of Applied Phycology*, 20(3), 245-251.
- Song, D., Fu, J., & Shi, D. (2008). Exploitation of oil-bearing microalgae for biodiesel. *Chinese Journal of Biotechnology*, 24(3), 341-348.
- Sparkman, O. D. (2000). Mass spectrometry desk reference 2. *Journal of the American Society for Mass Spectrometry*, 12(11), 1144.
- Spolaore, P., Joannis-Cassan, C., Duran, E., & Isambert, A. (2006). Commercial applications of microalgae. *Journal of bioscience and bioengineering*, 101(2), 87-96.
- Stansell, G. R., Gray, V. M., & Sym, S. D. (2012). Microalgal fatty acid composition: implications for biodiesel quality. *Journal of Applied Phycology*, 24(4), 791-801.
- Stern, D. (2009). *The Chlamydomonas sourcebook: organellar and metabolic processes* (Vol. 2): Academic Press.
- Strauss, H. (2004). 4 Ga of seawater evolution: Evidence from the sulfur isotopic composition of sulfate. *Sulfur biogeochemistry: past and present*, 379, 195.
- Štrojsová, A., & Dyhrman, S. T. (2008). Cell-specific β -N-acetylglucosaminidase activity in cultures and field populations of eukaryotic marine phytoplankton. *FEMS microbiology ecology*, 64(3), 351-361.
- Subashchandrabose, S. R., Ramakrishnan, B., Megharaj, M., Venkateswarlu, K., & Naidu, R. (2013). Mixotrophic cyanobacteria and microalgae as distinctive biological agents for organic pollutant degradation. *Environment international*, 51, 59-72.
- Suh, I. S., & Lee, C.-G. (2003). Photobioreactor engineering: design and performance. *Biotechnology and Bioprocess Engineering*, 8(6), 313.
- Surriya, O., Saleem, S. S., Waqar, K., Kazi, A. G., & Öztürk, M. (2015). Bio-fuels: a blessing in disguise *Phytoremediation for Green Energy* (pp. 11-54): Springer.

- Sushchik, N., Kalacheva, G., Zhila, N., Gladyshev, M., & Volova, T. (2003). A temperature dependence of the intra-and extracellular fatty-acid composition of green algae and cyanobacterium. *Russian journal of plant physiology*, 50(3), 374-380.
- Takeda, T., Nakano, Y., Takahashi, M., Konno, N., Sakamoto, Y., Arashida, R., . . . Suzuki, K. (2015). Identification and enzymatic characterization of an endo-1, 3- β -glucanase from *Euglena gracilis*. *Phytochemistry*, 116, 21-27.
- Thompson Jr, G. A. (1996). Lipids and membrane function in green algae. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism*, 1302(1), 17-45.
- Tibbetts, S. M., Milley, J. E., & Lall, S. P. (2015). Chemical composition and nutritional properties of freshwater and marine microalgal biomass cultured in photobioreactors. *Journal of Applied Phycology*, 27(3), 1109-1119.
- Touloupakis, E., Cicchi, B., Benavides, A. M. S., & Torzillo, G. (2016). Effect of high pH on growth of *Synechocystis* sp. PCC 6803 cultures and their contamination by golden algae (*Poterioochromonas* sp.). *Applied microbiology and biotechnology*, 100(3), 1333-1341.
- Trabelsi, L., Chaieb, O., Mnari, A., Abid-Essafi, S., & Aleya, L. (2016). Partial characterization and antioxidant and antiproliferative activities of the aqueous extracellular polysaccharides from the thermophilic microalgae *Graesiella* sp. *BMC complementary and alternative medicine*, 16(1), 210.
- Troncoso-Ponce, M. A., Garcés, R., & Martínez-Force, E. (2010). Glycolytic enzymatic activities in developing seeds involved in the differences between standard and low oil content sunflowers (*Helianthus annuus* L.). *Plant Physiology and Biochemistry*, 48(12), 961-965.
- Tsuzuki, M., Ohnuma, E., Sato, N., Takaku, T., & Kawaguchi, A. (1990). Effects of CO₂ concentration during growth on fatty acid composition in microalgae. *Plant physiology*, 93(3), 851-856.
- Tyson, R. (2012). *Sedimentary organic matter: organic facies and palynofacies*: Springer Science & Business Media.
- Tzovenis, I., De Pauw, N., & Sorgeloos, P. (2003). Optimisation of T-ISO biomass production rich in essential fatty acids: II. Effect of different light regimes on the production of fatty acids. *Aquaculture*, 216(1-4), 223-242.

- Van De Meene, A. M., Hohmann-Marriott, M. F., Vermaas, W. F., & Roberson, R. W. (2006). The three-dimensional structure of the cyanobacterium *Synechocystis* sp. PCC 6803. *Archives of Microbiology*, 184(5), 259-270.
- Van Der Straeten, D., Rodrigues-Pousada, R. A., Goodman, H. M., & Van Montagu, M. (1991). Plant enolase: gene structure, expression, and evolution. *The Plant Cell*, 3(7), 719-735.
- Van Wychen, S., & Laurens, L. M. (2013). *Determination of Total Carbohydrates in Algal Biomass: Laboratory Analytical Procedure (lap)*. Retrieved from
- Villas-Bôas, S., Roessner, U., Hansen, M., Smedsgaard, J., & Nielsen, J. Metabolome Analysis: An Introduction. 2007: John Wiley & Sons, Inc.
- Vitova, M., Bisova, K., Kawano, S., & Zachleder, V. (2015). Accumulation of energy reserves in algae: from cell cycles to biotechnological applications. *Biotechnology advances*, 33(6), 1204-1218.
- Voll, L. M., Hajirezaei, M. R., Czogalla-Peter, C., Lein, W., Stitt, M., Sonnewald, U., & Börnke, F. (2009). Antisense inhibition of enolase strongly limits the metabolism of aromatic amino acids, but has only minor effects on respiration in leaves of transgenic tobacco plants. *New Phytologist*, 184(3), 607-618.
- Vyverman, W. (2004). Experimental studies on sexual reproduction in diatoms. *International Review of Cytology: A Survey of Cell Biology*, 237, 91.
- Waghmare, A. G., Salve, M. K., LeBlanc, J. G., & Arya, S. S. (2016). Concentration and characterization of microalgae proteins from *Chlorella pyrenoidosa*. *Bioresources and Bioprocessing*, 3(1), 16.
- Wahidin, S., Idris, A., & Shaleh, S. R. M. (2013). The influence of light intensity and photoperiod on the growth and lipid content of microalgae *Nannochloropsis* sp. *Bioresource technology*, 129, 7-11.
- Wang, B., Wang, J., Zhang, W., & Meldrum, D. R. (2012). Application of synthetic biology in cyanobacteria and algae. *Frontiers in microbiology*, 3, 344.
- Wang, S.-B., Hu, Q., Sommerfeld, M., & Chen, F. (2003). An optimized protocol for isolation of soluble proteins from microalgae for two-dimensional gel electrophoresis analysis. *Journal of Applied Phycology*, 15(6), 485-496.
- Wang, S., Jiang, X., Wang, N., Yu, L., Li, Z., & He, P. (2007). Research on pyrolysis characteristics of seaweed. *Energy & Fuels*, 21(6), 3723-3729.

- Wang, W., Vignani, R., Scali, M., & Cresti, M. (2006). A universal and rapid protocol for protein extraction from recalcitrant plant tissues for proteomic analysis. *Electrophoresis*, 27(13), 2782-2786.
- Wang, X., Liu, X., & Wang, G. (2011). Two-stage Hydrolysis of Invasive Algal Feedstock for Ethanol Fermentation. *Journal of integrative plant biology*, 53(3), 246-252.
- Wang, X., Sheng, L., & Yang, X. (2017). Pyrolysis characteristics and pathways of protein, lipid and carbohydrate isolated from microalgae *Nannochloropsis* sp. *Bioresource technology*, 229, 119-125.
- Wang, Z. T., Ullrich, N., Joo, S., Waffenschmidt, S., & Goodenough, U. (2009). Algal lipid bodies: stress induction, purification, and biochemical characterization in wild-type and starchless *Chlamydomonas reinhardtii*. *Eukaryotic cell*, 8(12), 1856-1868.
- Wehr, J. D., Sheath, R. G., & Kociolek, J. P. (2015). *Freshwater algae of North America: ecology and classification*: Elsevier.
- Wei, L., Huang, X., & Huang, Z. (2015). Temperature effects on lipid properties of microalgae *Tetraselmis subcordiformis* and *Nannochloropsis oculata* as biofuel resources. *Chinese journal of oceanology and limnology*, 33(1), 99-106.
- Weignerová, L., Simerská, P., & Křen, V. (2009). α -Galactosidases and their applications in biotransformations. *Biocatalysis and Biotransformation*, 27(2), 79-89.
- Weiss, T. L., Roth, R., Goodson, C., Vitha, S., Black, I., Azadi, P., . . . Goodenough, U. (2012). Colony organization in the green alga *Botryococcus braunii* (Race B) is specified by a complex extracellular matrix. *Eukaryotic cell*, EC. 00184-00112.
- Wichelecki, D. J., Balthazor, B. M., Chau, A. C., Vetting, M. W., Fedorov, A. A., Fedorov, E. V., Hillerich, B. S. (2014). Discovery of function in the enolase superfamily: D-mannonate and D-gluconate dehydratases in the D-mannonate dehydratase subgroup. *Biochemistry*, 53(16), 2722-2731.
- Wilson, R. (2013). Sensitivity and specificity: twin goals of proteomics assays. Can they be combined? *Expert review of proteomics*, 10(2), 135-149.
- Winck, F. V., Melo, D. O. P., & Barrios, A. F. G. (2013). Carbon acquisition and accumulation in microalgae *Chlamydomonas*: Insights from “omics” approaches. *Journal of proteomics*, 94, 207-218.

- Wishart, D. S. (2016). Emerging applications of metabolomics in drug discovery and precision medicine. *Nature Reviews Drug Discovery*, 15(7), 473.
- Wold, F. (1971). 18 Enolase *The enzymes* (Vol. 5, pp. 499-538): Elsevier.
- Wolf, F. R., Nonomura, A. M., & Bassham, J. A. (1985). Growth and branched hydrocarbon production in a strain of *Botryococcus braunii* (chlorophyta) 1. *Journal of phycology*, 21(3), 388-396.
- Xiao, R., & Zheng, Y. (2016). Overview of microalgal extracellular polymeric substances (EPS) and their applications. *Biotechnology advances*, 34(7), 1225-1244.
- Xin, L., Hong-ying, H., Ke, G., & Ying-xue, S. (2010). Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalga *Scenedesmus* sp. *Bioresource technology*, 101(14), 5494-5500.
- Xu, H., Miao, X., & Wu, Q. (2006). High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters. *Journal of biotechnology*, 126(4), 499-507.
- Yamada, T., & Sakaguchi, K. (1982). Comparative studies on *Chlorella* cell walls: Induction of protoplast formation. *Archives of Microbiology*, 132(1), 10-13.
- Yamaguchi, K., Nakano, H., Murakami, M., Konosu, S., Nakayama, O., Kanda, M., . . . Iwamoto, H. (1987). Lipid composition of a green alga, *Botryococcus braunii*. *Agricultural and biological chemistry*, 51(2), 493-498.
- Yeesang, C., & Cheirsilp, B. (2011). Effect of nitrogen, salt, and iron content in the growth medium and light intensity on lipid production by microalgae isolated from freshwater sources in Thailand. *Bioresource technology*, 102(3), 3034-3040.
- Yin, D., Geng, Y., Mei, H., Ouyang, Z., Hu, H., & Li, Y. (2008). The effects of several environmental factors on the photosynthesis of *Botryococcus braunii*. *Journal of Wuhan Botanical Research*, 26(1), 64-69.
- Yokoyama, S., Hirota, H., Kigawa, T., Yabuki, T., Shirouzu, M., Terada, T., . . . Nishimura, Y. (2000). Structural genomics projects in Japan. *Nature Structural & Molecular Biology*, 7(11s), 943.
- Yoshimura, T., Okada, S., & Honda, M. (2013). Culture of the hydrocarbon producing microalga *Botryococcus braunii* strain Showa: Optimal CO₂, salinity, temperature, and irradiance conditions. *Bioresource technology*, 133, 232-239.

- Zhang, F., Kabeya, H., Kitagawa, R., Hirotsu, T., Yamashita, M., & Otsuki, T. (2000). A novel polyethylene–chlorella composite. I. Characterization of chlorella biologically fixing CO₂. *Journal of applied polymer science*, 77(10), 2278-2284.
- Zhang, J., & Hu, B. (2012). A novel method to harvest microalgae via co-culture of filamentous fungi to form cell pellets. *Bioresource technology*, 114, 529-535.
- Zhou, W., Min, M., Hu, B., Ma, X., Liu, Y., Wang, Q., Ruan, R. (2013). Filamentous fungi assisted bio-flocculation: a novel alternative technique for harvesting heterotrophic and autotrophic microalgal cells. *Separation and Purification Technology*, 107, 158-165.
- Zhu, L. (2015). Microalgal culture strategies for biofuel production: a review. *Biofuels, Bioproducts and Biorefining*, 9(6), 801-814.
- Zhu, L., Li, Z., & Hiltunen, E. (2016). Strategies for lipid production improvement in microalgae as a biodiesel feedstock. *BioMed research international*, 2016.
- Zhu, L., Wang, Z., Shu, Q., Takala, J., Hiltunen, E., Feng, P., & Yuan, Z. (2013). Nutrient removal and biodiesel production by integration of freshwater algae cultivation with piggery wastewater treatment. *Water research*, 47(13), 4294-4302.
- Zhu, S., Wang, Y., Huang, W., Xu, J., Wang, Z., Xu, J., & Yuan, Z. (2014). Enhanced accumulation of carbohydrate and starch in *Chlorella zofingiensis* induced by nitrogen starvation. *Applied biochemistry and biotechnology*, 174(7), 2435-2445.

